

FORMATIVE PROCESSES WITH APPLICATIONS TO
THE DECISION PROBLEM IN SET THEORY: II.
POWERSET AND SINGLETON OPERATORS,
FINITENESS PREDICATE*

Domenico Cantone (corresponding author)

Dipartimento di Matematica e Informatica, Università di Catania

Viale Andrea Doria 6, I-95125 Catania, Italy.

E-mail: cantone@dmf.unict.it Phone: +39 095 7383052 Fax: +39 095 330094

Pietro Ursino

Dipartimento di Matematica e Informatica, Università di Catania

E-mail: ursino@dmf.unict.it

*This research has been partially funded by PRIN project 2006/2007 ‘Large-scale development of certified mathematical proofs’.

Abstract

In this paper we solve the satisfiability problem for a quantifier-free fragment of set theory involving the powerset and the singleton operators and a finiteness predicate, besides the basic Boolean set operators of union, intersection, and difference. The more restricted fragment obtained by dropping the finiteness predicate has been shown to have a solvable satisfiability problem in a previous paper, by establishing for it a small model property. To deal with the finiteness predicate we have formulated and proved a *small model witness property* for our fragment of set theory, namely a property which asserts that any satisfiable formula of our fragment has a model admitting a “small” representation.

Key words Satisfiability decision problem, satisfaction algorithm, Zermelo-Fraenkel set theory.

1 Introduction

Since the late seventies, the decision problem in set theory has been investigated very actively, as part of a project aiming at the mechanical formalization of mathematics using a proof verifier based on the set-theoretic formalism [OS02, COS*03, OCP*06].

The large body of decidability results gave rise to the field of *Computable Set Theory*; the reader is referred to [CFO89, COP01] for a very comprehensive account.

Using specialized techniques, several fragments of set theory were shown to have a solvable satisfiability decision problem. But it soon became clear that in order to combine such decidable fragments into more comprehensive ones, more sophisticated methods were needed, which would make the growing mass of technicalities involved more manageable.

Such methods began to emerge in [Can91] and gained perspicuity and uniformity of approach in [CU97, COU02]; they rely on a careful analysis of the formation process of families of pairwise disjoint sets. This analysis brought into light decidable conditions which are necessary and sufficient to ensure satisfiability within a collection of set-theoretic formulae denoted **MLSSP** (viz., “*Multi-Level Syllogistic with Singleton and Powerset operators*”). Specifically, **MLSSP** is the quantifier-free language of set theory, consisting of all propositional combinations of literals involving the set operators of union, intersection, difference, powerset, finite enumerations, and the set relators of equality, membership, and inclusion.

Let us recall that the satisfiability problem for **MLSSP** was shown to be decidable by singling out a computable function c which associates a positive integer with every formula of **MLSSP** so that the following “small model property” holds: any satisfiable **MLSSP**-formula Φ can be satisfied by a set assignment mapping the variables of Φ into sets of finite rank bounded by $c(\Phi)$. Decidability then follows readily, in view of the facts that finitely many such assignments exist, that their collection can be effectively generated, and, additionally, for every such assignment one can effectively test whether or not it satisfies Φ . As an immediate by-product, one has that any satisfiable **MLSSP**-formula is satisfied by an assignment of finite sets (of bounded rank) to the variables.

A natural question then arises: how can one tackle the satisfiability problem regarding a collection of formulae which extends **MLSSP** and comprises formulae satisfiable only by means of assignments some of whose images are *infinite* sets?

The present paper contributes to answering this question by providing a positive solution to the satisfiability problem for the language **MLSSPF** which extends **MLSSP** by a predicate $Finite(x)$ asserting that its argument x has finite cardinality. Obviously, since the negated predicate $\neg Finite(x)$ expresses that x is an infinite set, **MLSSPF** cannot enjoy the above-described small model property; but, on the other hand, we will show that **MLSSPF** enjoys a less demanding property, called “small model witness property” (cf. [Urs05]), asserting the existence of a computable function c defined over the formulae of **MLSSPF** and such that any satisfiable **MLSSPF**-formula Φ has a “witness satisfying assignment” of finite rank bounded by $c(\Phi)$. Though a witness satisfying assignment for a formula Φ is not required to directly satisfy Φ , it represents a satisfying assignment for Φ through a “pumping mechanism” which will be illustrated at length in the paper. Additionally, since one can test algorithmically whether a given assignment of finite rank is a witness satisfying assignment for a formula Φ , decidability of the satisfiability problem for **MLSSPF** follows.

To achieve these results we will rely on the notion of formative process introduced in [COU02] and [COP01], which enables one to characterize any given collection of sets by means of a (transfinite) sequence of construction steps. By suitable manipulations,

namely through the pumping technique mentioned above, a formative process for a witness satisfying assignment can be turned into a formative process for a(n infinite) satisfying assignment.

We expect that similar techniques will demonstrate very useful in the investigation of the satisfiability problem for various fragments of set theory involving operators such as the unary union $\bigcup$ and the Cartesian product $\times$, since these can be used to express satisfiable formulae such as

$$x \subseteq \bigcup x \wedge x \neq \emptyset \quad \text{and} \quad x \times x \subseteq x \wedge x \neq \emptyset,$$

which admit only infinite satisfying assignments.

The paper is organized as follows. In Section 2, after reviewing some basic set-theoretic notions and terminology, we give a precise definition of the language **MLSSPF** of our interest and briefly discuss its satisfiability problem. Then in Section 3 we present and extend the notion of $\mathcal{P}$ -boards and the technique of formative processes. We also introduce the notion of *pumping path* and prove a related *pumping theorem* which is then used in the subsequent Section 4 as a basic tool for showing how to produce a satisfying assignment from a witness satisfying assignment. Section 5 contains some closing remarks and hints at possible directions of future research. Finally, an appendix containing the proof of a technical result used in Section 4 concludes the paper.

2 The satisfiability problem for **MLSSPF**

Before giving the precise definition of the language **MLSSPF** and discussing its satisfiability problem, we review some basic set-theoretic concepts and terminology.

2.1 Basic notions

Our considerations will refer to the von Neumann's cumulative hierarchy $\mathcal{V}$, often regarded as the intended model of Zermelo-Fraenkel set theory, also referred to as “{ sets }” throughout the paper, which is defined in the following recursive manner:

$$\begin{aligned} \mathcal{V}_0 &= \emptyset, \\ \mathcal{V}_{\alpha+1} &= \mathcal{P}(\mathcal{V}_\alpha), \quad \text{for each ordinal } \alpha, \\ \mathcal{V}_\lambda &= \bigcup_{\mu < \lambda} \mathcal{V}_\mu, \quad \text{for each limit ordinal } \lambda, \\ \mathcal{V} &= \bigcup_{\mu \in On} \mathcal{V}_\mu \end{aligned}$$

(where $\mathcal{P}(S)$ denotes the powerset of S and On is the class of all ordinal numbers — recall that a **LIMIT** ordinal is an ordinal which differs from 0 and from any **SUCCESSOR** ordinal, of the form $\alpha + 1$). Then we can readily define a rank function rk on $\mathcal{V}$ by putting

$$\text{rk}(S) \stackrel{\text{Def}}{=} \min\{\alpha \in On \mid S \in \mathcal{V}_{\alpha+1}\},$$

for each $S \in \mathcal{V}$ (see [Jec78] for further details).

A set S is **TRANSITIVE** if $\bigcup S \subseteq S$, i.e., every member of a member of S belongs to S . The **TRANSITIVE CLOSURE** $\text{TrCl}(S)$ of a set S is the minimal transitive set containing S , i.e.,

$$\text{TrCl}(S) \stackrel{\text{Def}}{=} \bigcap \{T \in \mathcal{V} \mid S \subseteq T \wedge T \text{ is transitive}\}.$$

A set S such that $\text{TrCl}(S)$ is finite is said to be HEREDITARILY FINITE. By ω we denote the rank of the collection of all hereditarily finite sets. It can be shown that ω is the smallest infinite (limit) ordinal.

We also recall that a partition Σ is a collection of pairwise disjoint nonnull sets, called BLOCKS of Σ . The union $\bigcup \Sigma$ of a partition Σ is the DOMAIN of Σ . If the domain of a partition Σ is transitive, then we will say that Σ is a TRANSITIVE PARTITION.

For the sake of completeness, we briefly review the notations which will be used in the rest of the paper. We denote: by X^Y , the set of all functions f from Y into X ; by $f[S]$, the multi-image $\{Y \mid (\exists z \in S) Y = f(z)\}$; by $R \circ S$, the map composition of R and S . To describe a function $f = \{\langle X, Y_X \rangle \mid X \in Z\}$, we will interchangeably use the notation $f = \{Y_X\}_{X \in Z}$ and the notation $X \xrightarrow{f} Y_X$ ($X \in Z$); moreover, when Z is an ordinal and accordingly f is a Z -sequence, we will make use of the notation $f = (Y_X)_{X < Z}$.

2.2 MLSSPF: assignments, models, and its satisfiability problem

The syntax of the quantifier-free language MLSSPF is defined as follows. The symbols of MLSSPF are:

- infinitely many variables $x, y, z, \dots$;
- the constant symbol $\emptyset$;
- the set operators $\cdot \cup \cdot, \cdot \cap \cdot, \cdot \setminus \cdot, \{\cdot, \dots, \cdot\}$, and $\mathcal{P}(\cdot)$;
- the set predicates $\cdot \subseteq \cdot, \cdot = \cdot, \cdot \in \cdot$, and $\text{Finite}(\cdot)$.

The set of MLSSPF-TERMS is the smallest collection satisfying the following conditions:

- all variables and the constant $\emptyset$ are MLSSPF-terms;
- if s and t are MLSSPF-terms, so are $s \cup t, s \cap t, s \setminus t$, and $\mathcal{P}(s)$;
- if $s_1, \dots, s_n$ are MLSSPF-terms, so is $\{s_1, \dots, s_n\}$.

MLSSPF-ATOMS have the form $s \subseteq t, s = t, s \in t$, and $\text{Finite}(s)$, where s, t are MLSSPF-terms.

MLSSPF-FORMULAE are propositional combinations of MLSSPF-atoms. MLSSPF-LITERALS are MLSSPF-atoms and their negations.

The semantics of MLSSPF is defined in the obvious way. A SET ASSIGNMENT M over a collection V of variables is any map from V into the universe $\mathcal{V}$ of all sets (in symbols, $M \in \mathcal{V}^V$ or $M \in \{\text{sets}\}^V$). The rank of a set assignment M over a collection V of variables is given by

$$\text{rk}(M) =_{\text{Def}} \max_{v \in V} \text{rk}(Mv).$$

If V is the collection of variables occurring in an MLSSPF-formula Φ , then a set assignment M over V is said to SATISFY Φ if the sentence Φ^M obtained by substituting in Φ each variable x by its interpretation Mx is true, where the operators and predicates in Φ are interpreted in the standard way. We will call any set assignment which satisfies a given MLSSPF-formula Φ a MODEL for Φ . When Φ has a model, we will say that Φ is SATISFIABLE.

The SATISFIABILITY PROBLEM for MLSSPF is the problem of establishing whether any given MLSSPF-formula is satisfiable.

By using a normalization procedure of the kind described in [CFO89, pp. 96–99], the satisfiability problem for MLSSPF can easily be reduced to the satisfiability problem for NORMALIZED CONJUNCTIONS of MLSSPF, namely conjunctions of literals of the following types:

$$\begin{array}{lllll} x = y, & x \neq y, & x = \emptyset, & x = y \cup z, & x = y \cap z, \\ x = y \setminus z, & x \subseteq y, & x \not\subseteq y, & x \in y, & x \notin y, \\ x = \mathcal{P}(y), & x = \{y_1, \dots, y_H\}, & \text{Finite}(y), & \neg \text{Finite}(y). \end{array}$$

By dropping from MLSSPF the literals involving the finiteness predicate, one obtains the theory MLSSP, whose decision problem has been solved in [Can91] and later revisited in [COU02] with the approach relying on formative processes.

2.3 Decision tests and Venn partitions

The decidability of MLSSP has been shown in [Can91, COU02] by proving that MLSSP enjoys a SMALL MODEL PROPERTY. This amounts to proving that there exists a computable function c defined over the collection of MLSSP-formulae such that any satisfiable MLSSP-formula Φ has a model of rank bounded by $c(\Phi)$. Therefore, to test the satisfiability of an MLSSP-formula Φ involving the variables V_Φ , one could execute the following procedure:

1. **for each** set assignment M over V_Φ , whose rank is limited by $c(\Phi)$ **do**
2. **if** M satisfies Φ **then**
3. **return** “ Φ is satisfiable”
4. **return** “ Φ is unsatisfiable”

The effectiveness of the above procedure follows immediately from the fact that there are only finitely many set assignments over V_Φ of finite bounded rank (and these can be effectively generated) and, additionally, it can be effectively verified whether any such set assignment satisfies the formula Φ .

In the case of MLSSPF, we will not be able to prove a small model property of the kind seen for MLSSP, since there are satisfiable MLSSPF-formulae which admit only models of infinite rank (e.g., $\neg \text{Finite}(x)$). Thus, we will only prove a weaker property for MLSSPF, the SMALL MODEL WITNESS PROPERTY, which still yields the decidability of the satisfiability problem for MLSSPF. To be more specific, we will show that there exists a computable function c' , defined over the the collection of MLSSPF-formulae, such that for any satisfiable MLSSPF-formula Φ there exists a (witness) set assignment M over the variables V_Φ of Φ , having finite rank bounded by $c'(\Phi)$, such that, by means of a “pumping process” to be defined later, M can be transformed into a set assignment M' which satisfies Φ . Since, as will be shown later, one can effectively test whether a set assignment over V_Φ of finite rank can be pumped into a model for Φ , the decidability of the satisfiability problem for MLSSPF will follow.

For the purpose of studying the satisfiability problem for MLSSPF, it is helpful to associate with any set assignment M over V its VENN PARTITION Σ_M , defined by

$$\Sigma_M =_{\text{def}} \left\{ \bigcap_{v \in U} M v \setminus \bigcup_{v \in V \setminus U} M v : U \subseteq V \mid U \neq \emptyset \right\} \setminus \{ \emptyset \},$$

i.e., the coarsest partition of $\bigcup_{v \in V} M v$ induced by the sets $M v$, for v in V .

It is also convenient to define the map $\mathfrak{S}_M : V \rightarrow \mathcal{P}(\Sigma_M)$, where

$$\mathfrak{S}_M(v) = \{\sigma \in \Sigma_M \mid \sigma \subseteq Mv\}, \text{ for } v \text{ in } V,$$

so that $Mv = \bigcup \mathfrak{S}_M(v)$ holds, for each v in V .

A basic technique to prove the decidability result of our interest consists in transforming a given model of an MLSSPF-formula Φ into a witness set assignment for Φ of bounded rank. Therefore, we need to find appropriate ways to manipulate set assignments.

A set assignment M over V can be manipulated through its associated Venn partition Σ_M as follows. Let Σ' be any partition of the same cardinality as Σ_M and let $\beta : \Sigma_M \rightarrow \Sigma'$ be a *bijection*. Then we define a set assignment M' over V by putting

$$M'v =_{\text{Def}} \bigcup \beta[\mathfrak{S}_M(v)], \text{ for } v \text{ in } V.$$

Plainly, the Venn partition of M' is just Σ' .

Next, let us assume that M satisfies an MLSSPF-normalized conjunction Φ .

Under which conditions on Σ' and β can we infer that the formula Φ is also satisfied by M' ?

If Φ involves only literals of the form

$$\begin{array}{llll} x = y, & x \neq y, & x = \emptyset, & x = y \cup z, \\ x = y \cap z, & x = y \setminus z, & x \subseteq y, & x \not\subseteq y, \end{array} \quad (1)$$

then it easily follows that M' satisfies Φ , without requiring any particular condition on Σ' and β .

If, in addition to literals of type (1), the conjunction Φ involves also literals of type

$$x \in y, \quad x \notin y, \quad x = \mathcal{P}(y), \quad x = \{y_1, \dots, y_H\}, \quad (2)$$

then in order for M' to satisfy Φ we require that the following three conditions hold (a proof of this fact can be found in [COU02, Lemma 10.1]):

Condition 1: Σ' *ϵ -simulates* Σ_M via β , i.e., $\bigcup \beta[X] \in \bigcup \beta[Y]$ if and only if $\bigcup X \in \bigcup Y$, for $X, Y \subseteq \Sigma_M$;

Condition 2: Σ' *$\mathcal{P}$ -simulates* Σ_M via β , i.e., $\bigcup \beta[X] = \mathcal{P}(\bigcup \beta[Y])$, if $\bigcup X = \mathcal{P}(\bigcup Y)$, for $X, Y \subseteq \Sigma_M$;

Condition 3: Σ' *L -simulates* Σ_M via β , i.e., $\bigcup \beta[X] = \{\bigcup \beta[Y_1], \dots, \bigcup \beta[Y_L]\}$, if $\bigcup X = \{\bigcup Y_1, \dots, \bigcup Y_L\}$, for $X, Y_1, \dots, Y_L \subseteq \Sigma_M$, where L is an upper bound for the value H in each literal of type $x = \{y_1, \dots, y_H\}$ in Φ .

If we further assume that besides literals of type (1) and (2) the formula Φ contains also literals of the form *Finite*(y) and $\neg$ *Finite*(y), then M' satisfies Φ if, in addition to Conditions 1–3, also the following condition holds:

Condition 4: $\beta(\sigma)$ is finite if and only if σ is finite, for $\sigma \in \Sigma_M$.

In fact, it is convenient to include among the Venn regions associated with M also an “outer” disjoint region $\mathcal{E}_M$, defined in such a way that $\Sigma_M \cup \{\mathcal{E}_M\}$ is a transitive partition whose domain is the transitive closure of the set $\{Mv : v \in V\}$, namely

$$\mathcal{E}_M = \text{TrCl}(\{Mv \mid v \in V\}) \setminus \bigcup \Sigma_M.$$

In the study of transitive partitions, we will make use of *$\mathcal{P}$ -boards* and *formative processes*, conceptual tools already introduced in [COP01, Chapter 9] and treated in greater depth, but in slightly different terms, in [COU02]. In the next section we review the definitions and basic facts related to $\mathcal{P}$ -boards and formative processes, suitably extended for the purpose of properly dealing with infinite blocks.

3 Syllogistic boards and formative processes

3.1 Basics on syllogistic boards

We consider a *finite* set $\mathcal{P}$, whose elements we will call **PLACES** and whose subsets we will call **NODES**. We assume that $\mathcal{P} \cap \mathcal{P}(\mathcal{P}) = \emptyset$, so that no node is a place, and vice versa. We will use these places and nodes as the vertices of a directed bipartite graph $\mathcal{G}$ of a special kind, which we will call a $\mathcal{P}$ -BOARD. The edges issuing from each place q are, mandatorily, all pairs q, B such that $q \in B \subseteq \mathcal{P}$. The remaining edges of $\mathcal{G}$ must lead from nodes to places; hence, $\mathcal{G}$ is fully characterized by the function

$$\mathcal{T} \in \mathcal{P}(\mathcal{P})^{\mathcal{P}(\mathcal{P})}$$

associating with each node A the set of all places t such that $\langle A, t \rangle$ is an edge of $\mathcal{G}$. The elements of $\mathcal{T}(A)$ are called the **TARGETS** of A . We will usually represent $\mathcal{G}$ simply by $\mathcal{T}$.

Places and nodes of a $\mathcal{P}$ -board are meant to represent the blocks σ , and the subsets Γ (or, quite often, their unionsets $\bigcup \Gamma$), of a transitive partition Σ , respectively. Thus, if we indicate by $q^{(\bullet)}$ the block which corresponds to the place q , and by $B^{(\bullet)}$ the set which corresponds to the node B , our understanding is that

- $q^{(\bullet)} \neq \emptyset$, for all $q \in \mathcal{P}$;
- $q^{(\bullet)} \cap p^{(\bullet)} = \emptyset$ when $q \neq p$, for all $q, p \in \mathcal{P}$;
- $B^{(\bullet)}$ is the sub-partition $\{q^{(\bullet)} : q \in B\}$ of the partition $\mathcal{P}^{(\bullet)} = \Sigma$, for all $B \subseteq \mathcal{P}$;
- every element of any element of a block in $\mathcal{P}^{(\bullet)}$, belongs to a block in $\mathcal{P}^{(\bullet)}$.

Moreover, the intended meaning of $t \in \mathcal{T}(A)$ is that $t^{(\bullet)} \cap \mathcal{P}^*(A^{(\bullet)}) \neq \emptyset$, where the following shorthand notation is used:

$$\mathcal{P}^*(X) \quad =_{\text{Def}} \quad \{v \subseteq \bigcup X \mid (\forall u \in X) u \cap v \neq \emptyset\}.$$

More precisely, *not only* $t^{(\bullet)} \cap \mathcal{P}^*(A^{(\bullet)}) \neq \emptyset$ *must hold when* $t \in \mathcal{T}(A)$, *but also* $s^{(\bullet)} \cap \mathcal{P}^*(A^{(\bullet)}) = \emptyset$ *when* $s \in \mathcal{P} \setminus \mathcal{T}(A)$.

Definition 1 A transitive partition Σ is said to **COMPLY WITH** $\mathcal{G}$ via $q \mapsto q^{(\bullet)}$, where $\mathcal{G}$ is a $\mathcal{P}$ -board, $q \mapsto q^{(\bullet)}$ belongs to $\Sigma^{\mathcal{P}}$ and $\mathcal{T}(A) = \{q \in \mathcal{P} \mid \mathcal{P}^*(A^{(\bullet)}) \cap q^{(\bullet)} \neq \emptyset\}$, if the latter function satisfies all the constraints associated with $\mathcal{G}$, as indicated above (in particular, this requires $q \mapsto q^{(\bullet)}$ to be injective). $\square$

Any such board is said to be *induced* by Σ (for short, a Σ -board). We denote a transitive Σ -board by a couple $(\Sigma, \mathcal{G})$, where Σ is a transitive partition and $\mathcal{G}$ is the induced $\mathcal{P}$ -board.

For the purposes of this paper, some additional structure must be superimposed on $\mathcal{P}$ -boards:

Definition 2 A $\mathcal{P}$ -board $\mathcal{G} = (\mathcal{T}, \mathcal{F}, \mathcal{Q})$ is said to be COLORED when it has

- a designated set $\mathcal{F}$ of places,
- a designated set $\mathcal{Q}$ of nodes, such that $D \in \mathcal{Q}$ holds whenever $D \subseteq B \in \mathcal{Q}$ (in short, $\bigcup \mathcal{P}[\mathcal{Q}] \subseteq \mathcal{Q}$), and
- a target function $\mathcal{T}$.

The places in $\mathcal{F}$ are said to be *red*, the ones in $\mathcal{P} \setminus \mathcal{F}$ are said to be *green*; the nodes in $\mathcal{Q}$ are called $\mathcal{P}$ -nodes. A node is red if all places in it are red, and green otherwise; a list of vertices is green if all vertices lying on it are green. $\square$

Assume that Φ belongs to MLSSP, and let $\mathcal{M} \in \{\text{sets}\}^{\mathcal{X}_\Phi}$ be a set-valued assignment defined on the collection $\mathcal{X}_\Phi$ of variables in Φ . We denote by $\Sigma_{\mathcal{X}_\Phi}$ the Venn partition of the set $\mathcal{M}[\mathcal{X}_\Phi]$, and by $\mathfrak{S}_{\mathcal{M}}$ the function $\mathfrak{S}_{\mathcal{M}} \in \mathcal{P}(\Sigma_{\mathcal{X}_\Phi})^{\mathcal{X}_\Phi}$ such that $\mathcal{M}(v) = \bigcup \mathfrak{S}_{\mathcal{M}}(v)$ holds for every v in $\mathcal{X}_\Phi$.

Remark 1 Observe that any formula Φ with variables $\mathcal{X}_\Phi$ of a language resulting from an extension of Multi Level Syllogistic can be modified, without affecting its satisfiability, in such a way any model $\mathcal{M}$ generates a transitive $\Sigma_{\mathcal{X}_\Phi}$ [Can91, pp.195-196]. Because of that, from now on we shall assume that $\Sigma_{\mathcal{X}_\Phi}$ is transitive, for any model $\mathcal{M}$ of a formula Φ with variables $\mathcal{X}_\Phi$.

The intended meaning of $r \in \mathcal{F}$ is that $|r^{(\bullet)}| < \aleph_0$; $B \in \mathcal{Q}$ means that $\mathcal{P}^*(B^{(\bullet)}) \subseteq \bigcup \mathcal{P}^{(\bullet)}$. To see that the latter constraint belongs to MLSSP, note that an alternative way of stating it is as $\mathcal{P}(\bigcup B^{(\bullet)}) \subseteq \bigcup \mathcal{P}^{(\bullet)}$ (this, globally, needs to be made explicit only for the inclusion-maximal nodes B in $\mathcal{Q}$). It is, in fact, obvious that $\mathcal{P}^*(B^{(\bullet)}) \subseteq \bigcup \mathcal{P}^{(\bullet)}$ follows from $\mathcal{P}(\bigcup B^{(\bullet)}) \subseteq \bigcup \mathcal{P}^{(\bullet)}$, because $\mathcal{P}^*(B^{(\bullet)}) \subseteq \mathcal{P}(\bigcup B^{(\bullet)})$; conversely, if B is a $\mathcal{P}$ -node, and the condition $\mathcal{P}^*(D^{(\bullet)}) \subseteq \bigcup \mathcal{P}^{(\bullet)}$ is satisfied by every $\mathcal{P}$ -node D (and hence it holds when $D \subseteq B$), then

$$\mathcal{P}(\bigcup B^{(\bullet)}) = \bigcup \mathcal{P}^*[\{D^{(\bullet)} : D \subseteq B\}] \subseteq \bigcup \mathcal{P}^*[\{D^{(\bullet)} : D \in \mathcal{Q}\}] \subseteq \bigcup \mathcal{P}^{(\bullet)}.$$

Likewise, the constraints $t^{(\bullet)} \cap \mathcal{P}^*(A^{(\bullet)}) \neq \emptyset$ associated with edges $\langle A, t \rangle$ (and also constraints of the form $s^{(\bullet)} \cap \mathcal{P}^*(A^{(\bullet)}) = \emptyset$) can be written in terms of the customary operator $\mathcal{P}$, because

$$\mathcal{P}^*(A^{(\bullet)}) = \mathcal{P}(\bigcup A^{(\bullet)}) \setminus \bigcup \{\mathcal{P}(\bigcup (A \setminus \{q\})^{(\bullet)}) : q \in A\}.$$

Observe, thirdly, that the transitivity condition $\bigcup \bigcup \mathcal{P}^{(\bullet)} \subseteq \bigcup \mathcal{P}^{(\bullet)}$ is reducible to the MLSSP fragment of the set-theoretic language, too, because it can be restated as $\bigcup \mathcal{P}^{(\bullet)} \subseteq \mathcal{P}(\bigcup \mathcal{P}^{(\bullet)})$.

In conclusion, we may view a colored board $\mathcal{G}$ as being the representation of a conjunction $\varphi \equiv \varphi_{\mathcal{G}}$ of constraints each of which either belongs to MLSSP or has the form $Finite(r)$. Here we are momentarily regarding each place q as the name of an unknown set; when the unknowns are replaced by concrete sets $q^{(\bullet)}$ within φ , the overall substitution must satisfy φ , and the sets $q^{(\bullet)}$ will then collectively form a partition Σ actually complying with φ . Let us now obtain $\widehat{\varphi} \equiv \widehat{\varphi}_{\mathcal{G}}$ by adding to φ all constraints $\neg Finite(g)$ with $g \in \mathcal{P} \setminus \mathcal{F}$. By investigating how we can get a solution to $\widehat{\varphi}$ out of a solution to φ (if possible), we will shed light on the conditions for the satisfiability of any formula of the quantifier-free language MLSSP extended with the *Finite* relator; and, ultimately, we will come to an algorithm which is able to test any formula of this extended language for satisfiability.

When affirmative literals of the form $Finite(v)$, with v in $\mathcal{X}_\Phi$, are also included in Φ , the $\Sigma_{\mathcal{X}_\Phi}$ -board can be colored naturally, by putting

- (a) $\mathcal{F} = \bigcup \{ \mathfrak{S}_{\mathcal{M}}(v) \mid \text{a literal of either the form } v = \{w_1, \dots, w_H\} \text{ or the form } \textit{Finite}(v) \text{ occurs in } \Phi \};$
- (b) $\mathcal{Q}$ is the minimal collection of nodes such that
- $\mathfrak{S}_{\mathcal{M}}(u) \in \mathcal{Q}$ for all literals of the form $u = \mathcal{P}(w)$ in Φ , and
 - $\bigcup \mathcal{P}[\mathcal{Q}] \subseteq \mathcal{Q}$.

This will be called THE CANONICAL BOARD OF THE ASSIGNMENT $\mathcal{M}$ FOR THE MLSSPF-FORMULA Φ .

3.2 Formative processes and basic events

We will now discuss a technique by which one can find a transitive partition

$$\Sigma_{\xi} = \{q^{(\bullet)} : q \in \mathcal{P}\}$$

complying with a given $\mathcal{P}$ -board by

- * constructing a ξ -sequence of partitions which is a *formative process* in the sense specified by Def. 3 below, by then
- * ascertaining that every edge $\langle A, t \rangle$ of the $\mathcal{P}$ -board has been *activated* along the process (in a sense to be clarified below), and by then
- * taking as Σ_{ξ} the last partition of the sequence.

Definition 3 Let Σ, Σ' be partitions such that

- every block $\sigma \in \Sigma$ has a block $\sigma' \in \Sigma'$ for which $\sigma \subseteq \sigma'$;
- $\sigma_0, \sigma_1 \subseteq \tau$ implies $\sigma_0 = \sigma_1$ when $\sigma_0, \sigma_1 \in \Sigma$ and $\tau \in \Sigma'$.

Then we say that Σ' FRAMES Σ .

When Σ' frames Σ , and moreover

$$\Sigma \neq \Sigma', \text{ and } \bigcup \Sigma' \setminus \bigcup \Sigma \subseteq \mathcal{P}^*(\Gamma) \text{ for some } \Gamma \subseteq \Sigma,$$

then the ordered pair Σ, Σ' is called an ACTION.

A (FORMATIVE) PROCESS is a sequence $(\Sigma_{\mu})_{\mu \leq \xi}$ of partitions, where ξ can be any ordinal and, for every ordinal $\nu < \xi$:

- $\Sigma_{\nu} = \emptyset$ if $\nu = 0$;
- $\Sigma_{\nu}, \Sigma_{\nu+1}$ is an action;
- Σ_{λ} frames Σ_{ν} , and $\bigcup \Sigma_{\lambda} \subseteq \bigcup \bigcup \{\Sigma_{\gamma} : \gamma \in \lambda\}$, for every limit ordinal λ such that $\nu < \lambda \leq \xi$.

For all $\mu \leq \xi$ and $\tau \in \Sigma_{\xi}$, we will designate by $\tau^{(\mu)}$ the unique set such that

$$\tau^{(\mu)} = \text{the } \sigma \in \Sigma_{\mu} \text{ such that } \sigma \subseteq \tau, \text{ if any exists, else } \emptyset.$$

A process $(\Sigma_{\mu})_{\mu \leq \xi}$ is said to be GREEDY if, for all $\nu < \xi$ and $\Gamma \subseteq \Sigma_{\nu}$, the following holds:

$$\mathcal{P}^*(\Gamma) \cap \left(\bigcup \Sigma_{\nu+1} \setminus \bigcup \Sigma_{\nu} \right) \neq \emptyset \text{ implies } \left(\mathcal{P}^*(\Gamma) \setminus \bigcup \Sigma_{\nu+1} \right) \cap \bigcup \Sigma_{\xi} = \emptyset.$$

If we take, along with a colored $\mathcal{P}$ -board $\mathcal{T}, \mathcal{F}, \mathcal{Q}$, a bijection $q \mapsto q^{(\bullet)}$ from the places $\mathcal{P}$ to the final partition Σ_ξ of a formative process, and if moreover Σ_ξ complies with $\mathcal{T}, \mathcal{F}, \mathcal{Q}$, we get what we will call a COLORED $\mathcal{P}$ -PROCESS: namely the quintuple $(\Sigma_\mu)_{\mu \leq \xi}, (\bullet), \mathcal{T}, \mathcal{F}, \mathcal{Q}$. Then, for all $\mu \leq \xi$, $\nu < \xi$, $p \in \mathcal{P}$, $B \subseteq \mathcal{P}$, we will designate by $p^{(\mu)}, B^{(\mu)}, B^{(\bullet)}, A_\nu$ the unique sets such that

$$\begin{aligned} p^{(\mu)} &= (p^{(\bullet)})^{(\mu)}; \\ \Delta^{(\nu)}(q) &= q^{(\nu+1)} \setminus \bigcup \mathcal{P}^{(\nu)}. \\ B^{(\mu)} &= \{q^{(\mu)} : q \in B\}, \text{ and } B^{(\bullet)} = \{q^{(\bullet)} : q \in B\}; \\ A_\nu &= \text{the set } A \subseteq \mathcal{P} \text{ for which } \bigcup \Sigma_{\nu+1} \setminus \bigcup \Sigma_\nu \subseteq \mathcal{P}^*(A^{(\nu)}). \end{aligned}$$

We will call A_ν the ν -TH MOVE of the process. $\square$

As an easy illustration of how one can reason about processes, let us prove the following:

Lemma 1 *In a process $(\Sigma_\mu)_{\mu \leq \xi}$, each Σ_μ frames Σ_ν , and $\bigcup \Sigma_\nu \subsetneq \bigcup \Sigma_\mu$, for all $\nu < \mu$.*

Proof. Arguing by contradiction, suppose that the statement is false, and let μ be the least ordinal such that, for some $\nu < \mu$, either $\bigcup \Sigma_\nu \subsetneq \bigcup \Sigma_\mu$ does not hold or Σ_ν is not framed by Σ_μ .

Trivially, since $\nu < \mu$, we have $\mu \neq 0$; moreover μ cannot be a successor ordinal $\mu = \gamma + 1$, else Σ_μ would frame Σ_γ , and Σ_γ would frame Σ_ν by the assumed minimality of μ , and hence, transitively, Σ_μ would frame Σ_ν , a contradiction.

But μ cannot be a limit ordinal either. In fact, if μ is a limit ordinal and we consider any ordinal $\nu < \mu$, then Σ_μ frames Σ_ν ; moreover, since $\nu + 1 \leq \mu$, we have $\bigcup \Sigma_\nu \subsetneq \bigcup \Sigma_{\nu+1} \subseteq \bigcup \Sigma_\mu$. $\blacksquare$

Relative to a formative process $(\Sigma_\mu)_{\mu \leq \xi}$, we can identify certain *events*: these, roughly speaking, are ordinals ν where something important is about to happen. Examples of events are, for any $\Gamma \subseteq \Sigma_\xi$ and $\sigma \in \Sigma_\xi$:

Edge-activation move: a $\nu < \xi$ such that $\sigma^{(\nu+1)} \cap \mathcal{P}^*(\Gamma) \neq \emptyset$ and $\sigma^{(\nu)} \cap \mathcal{P}^*(\Gamma) = \emptyset$;

Grand move: a $\nu < \xi$ such that $\bigcup \Gamma \in \bigcup \Sigma_{\nu+1} \setminus \bigcup \Sigma_\nu$.

Subtler events, which will be discussed later, create conditions that enable expansion of the block associated with a green place into an infinite block.

3.3 Historical profile of an infinite block

Let the triple $\mathcal{T}, \mathcal{F}, \mathcal{Q}$ represent a colored $\mathcal{P}$ -board: what guarantee do we have that a colored process $(\Sigma_\mu)_{\mu \leq \xi}, (\bullet), \mathcal{T}, \mathcal{F}, \mathcal{Q}$ exists? *A priori*, according to our definitions, none; as a matter of fact, should some place have no afferent edges, that is, should there be some $s \in \mathcal{P}$ such that $s \notin \mathcal{T}(A)$ for any $A \subseteq \mathcal{P}$, then no move A_ν would ever be allowed to bring elements into $s^{(\nu+1)}$, and consequently $s^{(\bullet)}$ could not be a block in a partition.

The situation is quite different if the colored $\mathcal{P}$ -board with which we start is induced by a transitive partition Σ , because then one can prove (cf. [COU02, p.176]) the following

Theorem 1 (Trace theorem) *If $\mathcal{T}, \mathcal{F}, \mathcal{Q}$ is a colored Σ -board induced by a finite transitive partition Σ , then a greedy colored process $(\Sigma_\mu)_{\mu \leq \xi}, \iota_\Sigma, \mathcal{T}, \mathcal{F}, \mathcal{Q}$ exists which ends with $\Sigma_\xi = \Sigma$ and has $|\xi| \leq |\bigcup \Sigma|$.* $\blacksquare$

As we showed in [COU02, p.180 ff.], if we have ‘in our pocket’, so to speak, this kind of oracular guarantee, namely a transitive partition Σ which induces a particular colored board, then we can imitate Σ by means of a ‘small’ partition Σ' , with rank

$$\text{rk}(\Sigma') \leq c(|\Sigma|),$$

where c is a specific computable function, namely $c(n) = \lceil \frac{25}{24} \rceil \cdot m + 5 \cdot 2^{2m+m} + 3$ with $m = \lceil \log_2(n+1) \rceil$. Instead of being obtained directly from Σ , this Σ' is obtained through a process $(\Sigma'_j)_{j \leq \ell}$ which ‘mimics’ a greedy colored process $(\Sigma_\mu)_{\mu \leq \xi}$ corresponding to Σ as stated in the trace theorem, and by then putting $\Sigma' = \Sigma'_\ell$. Here $\ell < \aleph_0$; moreover one can manage that the sequence of edge-activation moves and grand moves are ‘synchronized’—in a sense to be clarified below—between the two processes. In what follows, we will refer to $(\Sigma_\mu)_{\mu \leq \xi}$ and to $(\Sigma'_j)_{j \leq \ell}$, respectively, when speaking of the *original process* and of the *shadow process*.

Under one major respect, however, the way in which the small partition Σ' portrays Σ is unfaithful to the original: all of its blocks have in fact finite cardinality, regardless of the cardinality of corresponding blocks in the original partition. To find a remedy to this, we will first detect the causes which can make a block in Σ infinite: such causes are to some extent static, viz. features of the Σ -board, and to some other extent dynamic, viz. ‘*pumping*’ events which activate certain special paths of the Σ -board during the original process. On the basis of a careful analysis of how infinity can enter into play in the original process, we will discover how to infinitely expand, again, the green blocks of our small partition, taking advantage of two facts:

- the board of the original and of the shadow process is the same; and
- we can synchronize with each pumping event ν of the original process a regular chain of actions that can safely be interpolated next to the move of the shadow process corresponding to the original ν -th move. (‘Safely’ means that the imitation, in the sense of Def. 17 below, is not disrupted by the extra moves.)

3.3.1 Pumping paths

The block at a place s cannot become infinite, during a formative process, without some afferent edge carrying to s infinitely many elements drawn from a cycle. Otherwise stated, to make it possible that $|s^{(\bullet)}| \geq \aleph_0$, the board of the process must have a path $C_0, q_0, C_1, q_1, \dots, C_n, q_n, C_0, t_0, D_1, t_1, \dots, D_{m+1}, t_{m+1}$ ending with $t_{m+1} = s$. This fact should be easy to grasp, but we will nevertheless provide a proof of it soon. Notice that we can insist w.l.o.g. that, besides the double occurrence of C_0 , the only possible repetitions of vertices are $q_0 = t_0$ and $q_0 = t_0, C_1 = D_1$. These considerations lead to the following two definitions, of which the former is preparatory to the subsequent, very important Def. 5.

Definition 4 In a $\mathcal{P}$ -board $\mathcal{G}$, a PATH is an ordered vertex list $W_1, \dots, W_k$ within which places and nodes so alternate that W_i, W_{i+1} is an edge of $\mathcal{G}$ for $i = 1, \dots, k-1$. A path is said to be SIMPLE if no place or node occurs twice in it (i.e., $W_i \neq W_j$ when $0 < i < j \leq k$ and $i \equiv j \pmod{2}$).

If

$$\begin{aligned} \mathcal{D}' &\equiv q_0, C_1, \dots, q_n, C_{n+1} && \text{and} \\ \mathcal{D}'' &\equiv t_0, D_1, \dots, t_{m-1}, D_m, t_m, \end{aligned}$$

with $n, m \geq 0$, are *simple* paths in $\mathcal{G}$ which meet the following conditions,




Figure 1: The three kinds of pumping path

- $\langle C_{n+1}, t_0 \rangle$ is an edge of $\mathcal{G}$,
- $\{q_0, \dots, q_n\} \cap \{t_0, \dots, t_m\} \subseteq \{q_0\} \cap \{t_0\}$,
- $\{C_1, \dots, C_{n+1}\} \cap \{D_1, \dots, D_{m+1}\} \subseteq \{C_1\} \cap \{D_1\}$,
- if $C_1 = D_1$, then $q_0 = t_0$,

then $\mathcal{D}''$ is said to be CONTIGUOUS to $\mathcal{D}'$.

A path $W'_1, \dots, W'_\ell$ in $\mathcal{G}$ is said to be ADJACENT to another path $W_1, \dots, W_k$ in $\mathcal{G}$ if W_k, W'_ℓ is an edge of $\mathcal{G}$. $\square$

Definition 5 In a colored $\mathcal{P}$ -board $\mathcal{G} = (\mathcal{T}, \mathcal{F}, \mathcal{Q})$, consider paths

$$\begin{aligned} \mathcal{C} &\equiv C_0, q_0, C_1, \dots, q_n, C_{n+1} && \text{and} \\ \mathcal{D} &\equiv q_{n+1}, C_{n+2}, q_{n+2}, \dots, C_{n+m+1}, q_{n+m+1} \end{aligned}$$

(where $n, m \geq 0$), devoid of red places and such that $C_{n+1} = C_0$ and $\mathcal{D}$ is contiguous to the path $q_0, C_1, \dots, q_n, C_{n+1}$. (Hence, in particular, $q_{n+m+1} \in \mathcal{T}(C_{n+m+1}) \setminus \mathcal{F}$, and $q_i \in \mathcal{T}(C_i) \cap C_{i+1} \setminus \mathcal{F}$ for $i = 0, 1, \dots, n+m$.)

Then $\mathcal{C}$ is said to be a PUMPING CYCLE, and the path resulting from the concatenation of $\mathcal{D}$ at the end of $\mathcal{C}$ is said to be a PUMPING PATH. (cf. Figure 1.) $\square$

In what follows, we will set up the ground for a theorem generalizing the following easy proposition (whose proof we delay until we can generalize the statement):

Lemma 2 *Relative to a colored $\mathcal{P}$ -process $(\Sigma_\mu)_{\mu \leq \xi}, (\bullet), \mathcal{T}, \mathcal{F}, \mathcal{Q}$, consider a place s such that $|s^{(\bullet)}| \geq \aleph_0$. Then s lies on a pumping path.* ■

In sight of a proposition generalizing this one, which will follow from two other lemmas, we give the following definition:

Definition 6 For any node B and place t of a colored process,

$$\begin{aligned} \text{leastInf}(B) &=_{\text{Def}} \begin{cases} \text{the least ordinal } \mu \text{ for which } |\mathcal{P}^*(B^{(\mu)})| \geq \aleph_0, & \text{if } |\bigcup B^{(\bullet)}| \geq \aleph_0, \\ 0 & \text{otherwise;} \end{cases} \\ \text{leastInf}(t) &=_{\text{Def}} \text{leastInf}(\{t\}). \end{aligned}$$

Note that $\text{leastInf}(B) \neq 0$ and $\text{leastInf}(t) \neq 0$ imply respectively B and t to be green, else the ending partition would not comply with the color of the board. Note also that

$$\text{leastInf}(t) = \begin{cases} \text{the least ordinal } \mu \text{ for which } |t^{(\mu)}| \geq \aleph_0, & \text{if } |t^{(\bullet)}| \geq \aleph_0, \\ 0 & \text{otherwise.} \end{cases}$$

□

Here are the two lemmas preliminary to Thm.2, which will generalize Lemma 2:

Lemma 3 *Relative to a greedy colored $\mathcal{P}$ -process $(\Sigma_\mu)_{\mu \leq \xi}, (\bullet), \mathcal{T}, \mathcal{F}, \mathcal{Q}$ whose sequence of moves is $(A_\nu)_{\nu < \xi}$, consider a place s_0 such that $\text{leastInf}(s_0) = \nu + 1$ for some ordinal ν . Then there exist a place s_{k+1} and a green path $\mathcal{D} \equiv s_{k+1}, B_k, s_k, \dots, B_1, s_1, B_0, s_0$ from s_{k+1} to s_0 , with $k \geq 0$, such that*

- (a) $\text{leastInf}(s_{k+1})$ is a limit ordinal λ ;
- (b) $\text{leastInf}(s_{i+1}) \leq \text{leastInf}(B_i) < \text{leastInf}(s_i)$, for $i = 0, 1, \dots, k$;
- (c) for every edge $\langle B_i, s_i \rangle$ lying on $\mathcal{D}$ and for every node H other than B_i , it holds that

$$\begin{aligned} |s_i^{(\text{leastInf}(s_i))} \cap \mathcal{P}^*(B_i^{(\text{leastInf}(s_i)-1)})| &\geq \aleph_0 \quad \text{and} \\ |s_i^{(\text{leastInf}(s_i))} \cap \mathcal{P}^*(H^{(\text{leastInf}(s_i)-1)})| &< \aleph_0. \end{aligned}$$

Proof. Arguing by contradiction, let us assume that ν_0 is the least ordinal such that an s_0 with $\text{leastInf}(s_0) = \nu_0 + 1$ exists violating the thesis. (Clearly s_0 must be green.) Then $|\mathcal{P}^*(A_{\nu_0}^{(\nu_0)})| \geq \aleph_0$, where A_{ν_0} is the ν_0 -th move, and hence there is a green place $s_1 \in A_{\nu_0}$ such that $0 < \text{leastInf}(s_1) \leq \text{leastInf}(A_{\nu_0}) \leq \nu_0$. It can easily be checked that the edge $\langle A_{\nu_0}, s_0 \rangle$ satisfies condition (c). Thus, if $\text{leastInf}(s_1)$ were a limit ordinal, our initial assumption would be violated, because we could take $k = 0$ and $\mathcal{D} \equiv s_1, A_{\nu_0}, s_0$. Hence $\text{leastInf}(s_1) = \gamma + 1$, and therefore, by the assumed minimality of ν_0 , a green path $\mathcal{D}'$ leading to s_1 and satisfying the conditions of the lemma can be found. But then, prolonging $\mathcal{D}'$ first with A_{ν_0} and next with s_0 would give us a $\mathcal{D}$ whose existence conflicts with our initial assumption. ■

Lemma 4 *Consider a greedy process over a $\mathcal{P}$ -board $\mathcal{G}$, and let λ be a limit ordinal. Let, moreover, the subgraph $\mathcal{G}_\lambda$ of $\mathcal{G}$ consist of all*

- places q such that the set

$$I_{\lambda,q} = \{\nu < \lambda \mid q^{(\nu+1)} \neq q^{(\nu)}\}$$

is cofinal in λ (i.e., $\bigcup I_{\lambda,q} = \lambda$);

- nodes B such that the set

$$I_{\lambda}^B = \{\nu < \lambda \mid A_{\nu} = B\}$$

is cofinal in λ ;

- edges $\langle B, q \rangle$ such that

$$I_{\lambda,q}^B = \{\nu < \lambda \mid A_{\nu} = B \wedge q^{(\nu+1)} \neq q^{(\nu)}\}$$

is cofinal in λ ;

- edges $\langle q, B \rangle$ such that $q \in B$.

Then every vertex W of $\mathcal{G}_{\lambda}$ (if any exists) has at least one predecessor in $\mathcal{G}_{\lambda}$.

Proof. If W is a place p of $\mathcal{G}_{\lambda}$ then, since

$$I_{\lambda,p} = \bigcup \{I_{\lambda,p}^B : B \subseteq \mathcal{P}\},$$

from the assumption $\bigcup I_{\lambda,p} = \lambda$ and the finiteness of $\mathcal{P}$ it follows that $\bigcup I_{\lambda,p}^B = \lambda = \bigcup I_{\lambda}^B$ for some B : hence B is an immediate predecessor of p in $\mathcal{G}_{\lambda}$.

On the other hand, if W is a node B of $\mathcal{G}_{\lambda}$ then, from the assumption $\bigcup I_{\lambda}^B = \lambda$ and from the greediness of the process it follows that $\bigcup \{I_{\lambda,q} : q \in B\} = \lambda$. Therefore $\bigcup I_{\lambda,q} = \lambda$ for some $q \in B$, in view of the finiteness of B , and hence q is an immediate predecessor of B in $\mathcal{G}_{\lambda}$. ■

We can now combine the preceding two lemmas into the following important result:

Theorem 2 (Pumping theorem) *Relative to a greedy colored $\mathcal{P}$ -process $(\Sigma_{\mu})_{\mu \leq \xi}, (\bullet), \mathcal{T}, \mathcal{F}, \mathcal{Q}$ whose sequence of moves is $(A_{\nu})_{\nu < \xi}$, consider a place s such that $|s^{(\bullet)}| \geq \aleph_0$. Then s is the last node of a pumping path which can be split into adjacent paths $\mathcal{C}, \mathcal{D}', \mathcal{D}''$ such that*

1. $\mathcal{C}$ is a cycle;
2. the path $\mathcal{C}, \mathcal{D}'$ ends with a place g whose $\text{leastInf}(g)$ is a limit ordinal λ ;
3. $0 < \text{leastInf}(q) \leq \lambda = \bigcup I_{\lambda,q}^B$ holds, where $I_{\lambda,q}^B =_{\text{def}} \{\nu < \lambda \mid A_{\nu} = B \wedge q^{(\nu+1)} \neq q^{(\nu)}\}$, for every edge $\langle B, q \rangle$ lying on $\mathcal{C}, \mathcal{D}'$;
4. $\lambda \leq \text{leastInf}(B) < \text{leastInf}(q)$, for every edge $\langle B, q \rangle$ lying on $\mathcal{D}''$;
5. $\text{leastInf}(q) \leq \text{leastInf}(B)$, for every edge $\langle q, B \rangle$ lying on $\mathcal{D}''$;
6. for every edge $\langle D, p \rangle$ lying on $\mathcal{D}''$ and for every node H other than D , it holds that

$$\begin{aligned} |p^{(\text{leastInf}(p))} \cap \mathcal{P}^*(D^{(\text{leastInf}(p)-1)})| &\geq \aleph_0 & \text{and} \\ |p^{(\text{leastInf}(p))} \cap \mathcal{P}^*(H^{(\text{leastInf}(p)-1)})| &< \aleph_0. \end{aligned}$$

Proof. If $\text{leastInf}(s)$ is a limit ordinal, then we can take $\lambda = \text{leastInf}(s)$, $g = s$, and $\mathcal{D}''$ void. To construct $\mathcal{C}, \mathcal{D}'$, we can exploit Lemma 4, where $\mathcal{G}_{\lambda}$ turns out to be non-void since it contains s . Thanks to the fact that every vertex of $\mathcal{G}_{\lambda}$ has a predecessor in $\mathcal{G}_{\lambda}$,

starting with s and proceeding in a backward fashion we will sooner or later hit the same vertex twice, since $\mathcal{G}_\lambda$ is finite.

If $\text{leastInf}(s)$ is a successor ordinal, then by Lemma 3 there exists a green path

$$\overline{\mathcal{D}''} \equiv s_{k+1}, B_k, s_k, \dots, B_1, s_1, B_0, s_0, \text{ with } s_0 = s,$$

such that

- $\text{leastInf}(s_{k+1})$ is a limit ordinal λ ;
- $\text{leastInf}(s_{i+1}) \leq \text{leastInf}(B_i) < \text{leastInf}(s_i)$, for $i = 0, 1, \dots, k$;
- for every edge $\langle B_i, s_i \rangle$ lying on $\overline{\mathcal{D}''}$ and for every node H other than B_i , it holds that

$$\begin{aligned} \left| s_i^{(\text{leastInf}(s_i))} \cap \mathcal{P}^*(B_i^{(\text{leastInf}(s_i)-1)}) \right| &\geq \aleph_0 & \text{and} \\ \left| s_i^{(\text{leastInf}(s_i))} \cap \mathcal{P}^*(H^{(\text{leastInf}(s_i)-1)}) \right| &< \aleph_0. \end{aligned}$$

Let us put $g \equiv_{\text{Def}} s_{k+1}$. To construct the paths $\mathcal{C}, \mathcal{D}', \mathcal{D}''$ of the theorem, we argue as follows. Much as in the previous case, we use again Lemma 4 to construct a pumping path $\mathcal{C}, \mathcal{D}'$ leading to s_{k+1} . Starting in $\mathcal{G}_\lambda$ with s_{k+1} , we proceed backwards until either we hit the same vertex twice, or we hit the vertex B_k , provided that B_k is in $\mathcal{G}_\lambda$. In the latter case, we form a cycle by making a further backward step through the edge $\langle s_{k+1}, B_k \rangle$. Finally, we put $\mathcal{D}'' \equiv_{\text{Def}} B_k, s_k, \dots, B_1, s_1, B_0, s_0$.

It is an easy matter to verify that the paths $\mathcal{C}, \mathcal{D}', \mathcal{D}''$ satisfy the conditions of the theorem. ■

3.3.2 Grand events and local trash

In what follows, we will refine into a useful proposition the following easy remark: the block at a place s belonging to a $\mathcal{P}$ -node A cannot become infinite during a colored process unless A has a green place among its targets. To see this, assume that $s \in A \in \mathcal{Q}$ and $|s^{(\bullet)}| \geq \aleph_0$. Consequently, $|\mathcal{P}^*(A^{(\bullet)})| > \aleph_0$ and $\mathcal{P}^*(A^{(\bullet)}) \subseteq \bigcup \mathcal{P}^{(\bullet)}$, and hence there must be a place g such that $|\mathcal{P}^*(A^{(\bullet)}) \cap g^{(\bullet)}| > \aleph_0$, because $|\mathcal{P}^{(\bullet)}| = |\mathcal{P}| < \aleph_0$; which obviously implies that $g \in \mathcal{T}(A) \setminus \mathcal{F}$.

In sight of generalizing the above remark, recalling the notion of grand move and noticing that such an event occurs, in a colored process, at most once for each node A , we give the following definition of *grand event* $\text{GE}(A)$ associated with A :

Definition 7 For every node A and every ν such that $0 \leq \nu < \xi$,

$$\text{GE}(A) \equiv_{\text{Def}} \begin{cases} \text{the ordinal } \nu \text{ for which } \bigcup A^{(\bullet)} \in \bigcup \mathcal{P}^{(\nu+1)} \setminus \bigcup \mathcal{P}^{(\nu)}, & \text{if any exists,} \\ \text{the length } \xi \text{ of the process} & \text{otherwise.} \end{cases}$$

Moreover, for any given collection $\mathcal{A}$ of nodes, we put

$$\text{GE}(\mathcal{A}) \equiv_{\text{Def}} \min\{\text{GE}(A) : A \in \mathcal{A}\}.$$

□

Notice that this definition implies that for any node A and any ν such that $0 \leq \nu < \xi$,

$$\begin{aligned} \nu \leq \text{GE}(A) &\leftrightarrow \bigcup A^{(\bullet)} \notin \bigcup \mathcal{P}^{(\nu)}, \\ \nu = \text{GE}(A) &\leftrightarrow \bigcup A^{(\bullet)} \in \bigcup \mathcal{P}^{(\nu+1)} \setminus \bigcup \mathcal{P}^{(\nu)}, \\ \nu > \text{GE}(A) &\leftrightarrow \bigcup A^{(\bullet)} \in \bigcup \mathcal{P}^{(\nu)}. \end{aligned}$$

Other elementary properties, whose proofs are left to the reader, are stated in the following lemma.

Lemma 5 *Let $(\Sigma_\mu)_{\mu \leq \xi}, (\bullet), \mathcal{T}, \mathcal{F}, \mathcal{Q}$ be a greedy colored $\mathcal{P}$ -process and let $A \subseteq \mathcal{P}$ be a node. Then*

- (a) $A^{(\alpha)} = A^{(\bullet)}$, where $\alpha = \text{GE}(A)$;
- (b) if $q^{(\nu+1)} \supsetneq q^{(\nu)}$, for some $q \in A$ and some $\nu < \xi$, then $\text{GE}(A) > \nu$.

Proof. Straightforward. ■

The following definitions show properties which allow one to “drain off” the “residual” elements of a pumping process:

Definition 8 A place g is said to be a **LOCAL TRASH** for a node A and a colored $\mathcal{P}$ -process $(\{q^{(\mu)}\}_{q \in \mathcal{P}})_{\mu \leq \xi}, (\bullet), \mathcal{T}, \mathcal{F}, \mathcal{Q}$ if

- $g \in \mathcal{T}(A) \setminus \mathcal{F}$, i.e., g is a green target of A ;
- every node B such that $g \in B$ has $\text{GE}(B) > \text{GE}(A)$. □

Lemma 6 *Relative to a greedy colored $\mathcal{P}$ -process $(\Sigma_\mu)_{\mu \leq \xi}, (\bullet), \mathcal{T}, \mathcal{F}, \mathcal{Q}$, consider a node $A \subseteq \mathcal{P}$ such that $\bigcup A^{(\bullet)}$ is infinite and $\alpha = \text{GE}(A) < \xi$. Then the set $\mathcal{P}^*(A^{(\alpha)}) \setminus \bigcup \mathcal{P}^{(\alpha)}$ is infinite.*

Proof. Let $(A_\nu)_{\nu < \xi}$ be the sequence of moves of the given $\mathcal{P}$ -process and let

$$u_A =_{\text{Def}} \bigcup A^{(\alpha)} \setminus \bigcup A^{(\nu_0)},$$

where $\nu_0 =_{\text{Def}} \min\{\nu \leq \alpha : A_\nu = A\}$. If u_A is finite then, since $A^{(\alpha)}$ is finite and $\bigcup A^{(\alpha)} = \bigcup A^{(\bullet)}$ is infinite, the set $\{v \in \mathcal{P}^*(A^{(\alpha)}) : u_A \subseteq v\}$ is an infinite subset of $\mathcal{P}^*(A^{(\alpha)})$ disjoint from $\bigcup \mathcal{P}^{(\alpha)}$, and therefore $\mathcal{P}^*(A^{(\alpha)}) \setminus \bigcup \mathcal{P}^{(\alpha)}$ is infinite.

On the other hand, if u_A is infinite, then the set $\{\bigcup A^{(\alpha)} \setminus \{w\} : w \in u_A\}$ is an infinite subset of $\mathcal{P}^*(A^{(\alpha)})$. Since $|\{\bigcup A^{(\alpha)} \setminus \{w\} : w \in u_A\} \cap \bigcup \mathcal{P}^{(\alpha)}| \leq 1$, we again conclude that $\mathcal{P}^*(A^{(\alpha)}) \setminus \bigcup \mathcal{P}^{(\alpha)}$ is infinite. ■

Lemma 7 *Relative to a greedy colored $\mathcal{P}$ -process $(\Sigma_\mu)_{\mu \leq \xi}, (\bullet), \mathcal{T}, \mathcal{F}, \mathcal{Q}$, consider a node $A \subseteq \mathcal{P}$ such that $\bigcup A^{(\bullet)}$ is infinite. Then A has a local trash p such that $p^{(\bullet)}$ is infinite.*

Proof. Let $\alpha = \text{GE}(A)$. Since A is a node, we have $\mathcal{P}^*(A^{(\alpha)}) = \mathcal{P}^*(A^{(\bullet)}) \subseteq \bigcup \mathcal{P}^{(\bullet)}$; hence $\mathcal{P}^*(A^{(\alpha)}) \setminus \bigcup \mathcal{P}^{(\alpha)} \subseteq \bigcup \mathcal{P}^{(\alpha+1)}$, by the greediness of the process. Moreover, since $\mathcal{P}^*(A^{(\alpha)}) \setminus \bigcup \mathcal{P}^{(\alpha)}$ is an infinite set by Lemma 6, there must be a place $p \in \mathcal{T}(A)$ for which the set $p^{(\alpha+1)} \cap (\mathcal{P}^*(A^{(\alpha)}) \setminus \bigcup \mathcal{P}^{(\alpha)})$ is infinite too. Hence $p^{(\bullet)}$ is infinite, and therefore $p \notin \mathcal{F}$.

Moreover, since $p^{(\alpha+1)} \setminus p^{(\alpha)} \neq \emptyset$, it readily follows that $\text{GE}(B) > \alpha = \text{GE}(A)$ must hold for every node $B \subseteq \mathcal{P}$ such that $p \in B$, which proves that p is a local trash for A . ■

Given a path $\mathcal{D}$ in a $\mathcal{P}$ -board $\mathcal{G}$, we denote by $(\mathcal{D})_{\text{places}}$ and $(\mathcal{D})_{\text{nodes}}$ the collections of places and of nodes occurring in $\mathcal{D}$, respectively. Moreover, given a node B in $\mathcal{G}$, we denote by $\mathcal{N}(B)$ the collection of all nodes which have nonnull intersection with B .

Definition 9 Let $\mathcal{C}, \mathcal{D}$ be a pumping path relative to a given colored $\mathcal{P}$ -process $(\Sigma_\mu)_{\mu \leq \ell}, [\bullet], \mathcal{T}, \mathcal{F}, \mathcal{Q}$, with ℓ finite. Then $\langle (\mathcal{C}, \overline{\mathcal{D}}_0, \mathcal{D}_1, \dots, \mathcal{D}_N), \vec{m}, q_0, \sigma \rangle$ is called a *pumping chain* if

- $\overline{\mathcal{D}}_0, \mathcal{D}_1, \dots, \mathcal{D}_N$ is a partition of $\mathcal{D}$ into contiguous paths each ending with a place, where in particular
- $\mathcal{D}_N$ ends with the place σ ;
- $\vec{m} = m_0, m_1, \dots, m_N$ is a non-decreasing sequence $m_0 \leq m_1 < m_2 < \dots < m_N < \xi$ of ordinals; and
- putting $\mathcal{D}_0 \equiv_{\text{def}} \mathcal{C}, \overline{\mathcal{D}}_0$, we have:
 - (i) $q_0 \in (\mathcal{C})_{\text{places}}$ and $q^{[m_0]} \setminus \bigcup \bigcup \mathcal{P}^{[m_0]} \neq \emptyset$;
 - (ii) $\text{GE}(\mathcal{N}((\mathcal{D}_i)_{\text{places}})) \geq m_i$, for $i = 0, 1, \dots, N$;
 - (iii) $\mathcal{P}^*(B^{[m_i]}) \neq \emptyset$ (i.e., $\emptyset \notin B^{[m_i]}$), for $B \in (\mathcal{D}_i)_{\text{nodes}}$ and $i = 0, 1, \dots, N$. □

Remark 2 For a wholesale re-use of all notation concerning formative processes, we designate the entities associated with the shadow process as $\hat{\mathcal{Z}}^{[i]}$ ($\hat{q}^{[i]}$, $\hat{B}^{[i]}$, $\Delta^{[i]}(q)$, etc.).

Related to a weak “shadow” formative process $\hat{\omega} = (\{\hat{q}^{[i]}\}_{q \in \mathcal{P}})_{i \leq \ell}$, we use the collection of “salient” steps $\{\mu_i\}_{i \leq \ell}$ of the original process, as defined in [COU02, Def. 7.1].

Definition 10 A set $\mathcal{W}$ of places is said to be CLOSED for a colored $\mathcal{P}$ -process $(\{q^{(\mu)}\}_{q \in \mathcal{P}})_{\mu \leq \xi}, (\bullet), \mathcal{T}, \mathcal{F}, \mathcal{Q}$ if

- all of its elements are green;
- every $\mathcal{P}$ -node which intersects $\mathcal{W}$ has a local trash belonging to $\mathcal{W}$.

If $\mathcal{C}$ is a path in a $\mathcal{P}$ -board, we define a collection of places B to be a CLOSURE of $\mathcal{C}$ when B is closed and it contains $(\mathcal{C})_{\text{places}}$. A path $\mathcal{C}$ is said to be CLOSURE EQUIPPED whenever it owns a closure. □

Remark 3 Observe that the following function $i \mapsto \mu_i$ is an order isomorphism between the following structures $(\ell, <)$ and $(\{\mu_i\}_{i \leq \ell}, <)$. This fact has valuable consequences:

- (i) For every node B , $\mu_{\text{GE}(\hat{B})} = \text{GE}(B)$;
- (ii) g is a local trash for A if and only if $\hat{g}$ is a local trash for $\hat{A}$;
- (iii) $\hat{\mathcal{C}}$ is closure equipped if and only if $\mathcal{C}$ is closure equipped.

Remark 4 The following properties are easily verified.

- $q^{[m_0]} \setminus q^{[m_0-1]} \neq \emptyset$ implies $q^{[m_0]} \setminus \bigcup \bigcup \mathcal{P}^{[m_0]} \neq \emptyset$.
- If $\mathcal{P}^*(B^{(\mu_j)}) \neq \emptyset$ then $\mathcal{P}^*(\hat{B}^{[j]}) \neq \emptyset$.

The following is the key result of the present section. Roughly speaking, it shows that the “shadow” formative process $\hat{\omega} = (\{\hat{q}^{[i]}\}_{q \in \mathcal{P}})_{i \leq \ell}$ discovered in [COU02] preserves the pumping chains.

Lemma 8 *If $\mathcal{P}$ with greedy formative process $\varpi = (\{q^{(\alpha)}\}_{q \in \mathcal{P}})_{\alpha \leq \xi}$ has an infinite place $p^{(\bullet)}$, then $\widehat{\mathcal{P}}$, with weak “shadow” formative process $\widehat{\varpi} = (\{\widehat{q}^{[i]}\}_{q \in \mathcal{P}})_{i \leq \ell}$ and with salient steps $\{\mu_i\}_{i \leq \ell}$, owns a pumping chain at $\widehat{p}$ which is closure equipped.*

Proof. By Thm.2, p lies on a pumping path which can be split into paths $\mathcal{C}, \mathcal{D}', \mathcal{D}''$ such that

1. $\mathcal{C}$ is a cycle;
2. the path $\mathcal{C}, \mathcal{D}'$ ends with a place g whose $\text{leastInf}(g)$ is a limit ordinal λ ;
3. $0 < \text{leastInf}(q) \leq \lambda = \bigcup I_{q,\lambda}^B$ holds, where

$$I_{q,\lambda}^B =_{\text{Def}} \{\nu < \lambda \mid A_\nu = B \wedge q^{(\nu+1)} \neq q^{(\nu)}\},$$

for every edge $\langle B, q \rangle$ lying on $\mathcal{C}, \mathcal{D}'$;

4. $\lambda \leq \text{leastInf}(B) < \text{leastInf}(q)$, for every edge $\langle B, q \rangle$ lying on $\mathcal{D}''$;
5. $\text{leastInf}(q) \leq \text{leastInf}(B)$, for every edge $\langle q, B \rangle$ lying on $\mathcal{D}''$.

We will show that $\mathcal{C}, \mathcal{D}', \mathcal{D}''$ is actually a pumping chain relative to the shadow formative process $\widehat{\varpi}$.

Let

$$\widehat{\lambda} =_{\text{Def}} \min\{i \mid \mu_i \geq \lambda\}.$$

Then we define

$$m_0 =_{\text{Def}} \max \{1 \leq i \leq \widehat{\lambda} \mid \langle A_{\mu_{i-1}}, q \rangle \text{ lies on } \mathcal{C} \text{ and } \widehat{q}^{[i]} \setminus \widehat{q}^{[i-1]} \neq \emptyset \text{ for some place } q\}.$$

Notice that the definition of m_0 is well-given, i.e.,

$$\{i \mid 1 \leq i \leq \widehat{\lambda}, \langle A_{\mu_{i-1}}, q \rangle \text{ lies on } \mathcal{C} \text{ and } \widehat{q}^{[i]} \setminus \widehat{q}^{[i-1]} \neq \emptyset \text{ for some place } q\} \neq \emptyset.$$

Indeed, let $q \in (\mathcal{C})_{\text{places}}$ and let B be the predecessor of q in $\mathcal{C}$. Let $\mu_{j_0} \in \text{Sal}$ be such that $B = A_{\mu_{j_0}}$, $q^{(\mu_{j_0})} \cap \mathcal{P}^*(B^{(\mu_{j_0})}) = \emptyset$, and $q^{(\mu_{j_0}+1)} \cap \mathcal{P}^*(B^{(\mu_{j_0})}) \neq \emptyset$. Since $I_{\lambda,q}^B \neq \emptyset$, we have $\mu_{j_0} < \lambda \leq \mu_{\widehat{\lambda}}$, so that $1 \leq j_0 + 1 \leq \widehat{\lambda}$. Therefore,

$$j_0 + 1 \in \{i \mid 1 \leq i \leq \widehat{\lambda}, \langle A_{\mu_{i-1}}, q \rangle \text{ lies on } \mathcal{C} \text{ and } \widehat{q}^{[i]} \setminus \widehat{q}^{[i-1]} \neq \emptyset \text{ for some place } q\}.$$

Next we define $\overline{\mathcal{D}}_0$ to be the longest initial subpath of $\mathcal{D}', \mathcal{D}''$ which contains only nodes occurring in $(\mathcal{C})_{\text{nodes}}$.

Assuming that we have already defined the paths $\overline{\mathcal{D}}_0, \mathcal{D}_1, \dots, \mathcal{D}_k$, each ending with a place, and the integers $m_0 \leq m_1 < \dots < m_k$, for some $k \geq 0$, we define $\mathcal{D}_{k+1}$ and m_{k+1} as follows, provided that $\mathcal{D}', \mathcal{D}'' \not\equiv \overline{\mathcal{D}}_0, \mathcal{D}_1, \dots, \mathcal{D}_k$. Let D_{k+1} be the first node in $\mathcal{D}', \mathcal{D}''$ not occurring in $\overline{\mathcal{D}}_0, \mathcal{D}_1, \dots, \mathcal{D}_k$. We define m_{k+1} as the least integer $j \geq m_k$ such that $\mathcal{P}^*(\widehat{D}_{k+1}^{[j]}) \neq \emptyset$. Moreover, we define $\mathcal{D}_{k+1}$ as the longest subpath $\mathcal{E}$ of $\mathcal{D}', \mathcal{D}''$ starting at D_{k+1} , ending at a place, and such that $\mathcal{P}^*(\widehat{D}^{[m_{k+1}]}) \neq \emptyset$, for each $D \in (\mathcal{E})_{\text{nodes}}$.

Since $\mathcal{D}', \mathcal{D}''$ is finite, the above definitional process must terminate and produce a finite sequence $m_0 \leq m_1 < \dots < m_N$ of natural numbers and a partition $\overline{\mathcal{D}}_0, \mathcal{D}_1, \dots, \mathcal{D}_N$ of $\mathcal{D}', \mathcal{D}''$ into contiguous paths each ending with a place and such that $\mathcal{D}_1$ starts with the first node in $\mathcal{D}', \mathcal{D}''$ not occurring in $\mathcal{C}$. Finally, we put $\mathcal{D}_0 \equiv_{\text{Def}} \mathcal{C}, \overline{\mathcal{D}}_0$.

To show that $\mathcal{C}, \mathcal{D}', \mathcal{D}''$ is a pumping chain for the shadow process, it is now enough to verify that conditions (i), (ii), and (iii) of Def. 9 are satisfied, relative to the above partition of $\mathcal{C}, \mathcal{D}', \mathcal{D}''$ and to the sequence of natural numbers $m_0 \leq m_1 < \dots < m_N$.

Concerning condition (i), it is enough to observe that by the very definition of m_0 we must have $q^{[m_0]} \setminus q^{[m_0-1]} \neq \emptyset$, for some $q \in (\mathcal{C})_{places}$, so that $q^{[m_0]} \setminus \bigcup \mathcal{P}^{[m_0]} \neq \emptyset$ follows immediately.

The proof that condition (ii) of Def. 9 is satisfied can be simplified by the following properties(1-6), whose proofs are straightforward and are therefore omitted.

Property 1 *Let $q \in \mathcal{P}$ and let $B \subseteq \mathcal{P}$ such that $q \in B$. Then $\text{GE}(B) \geq \text{leastInf}(q)$.*

Property 2 *Let $q \in (\mathcal{D}'')_{places}$. Then*

(a) $\text{leastInf}(q) > \lambda$;

(b) *for each node $D \in (\mathcal{D}'')_{nodes}$ which precedes q in $\mathcal{D}''$, we have $\text{leastInf}(q) > \text{leastInf}(D)$.*

Proof. The property follows immediately from conditions 4 and 5 of Thm.2. ■

Property 3 *Let $i \in \{1, \dots, N\}$ and assume also that $m_{i-1} < m_i$. If D_i is the initial node of $\mathcal{D}_i$, then*

$$\min\{\mu < \xi \mid \mathcal{P}^*(D_i^{(\mu)}) \neq \emptyset\} = \mu_{m_{i-1}} + 1.$$

Property 4 *Let $D \in (\mathcal{C}, \mathcal{D}')_{nodes}$. Then*

$$\min\{\mu < \xi \mid \mathcal{P}^*(D^{(\mu)}) \neq \emptyset\} < \lambda.$$

Property 5 *Let $q \in (\mathcal{C}, \mathcal{D}')_{places}$ and let $B \subseteq \mathcal{P}$ be a node such that $q \in B$. Then $\text{GE}(A) \geq \lambda$.*

Property 6 *Let $i \in \{0, 1, \dots, \hat{\lambda}\}$. Then $\mu_i < \lambda$.*

To show that condition (ii) of Def. 9 holds, it is enough to prove that for every index $i \in \{0, 1, \dots, N\}$, place $q \in (\mathcal{D}_i)_{places}$, and node $B \subseteq \mathcal{P}$ such that $q \in B$, we have $\text{GE}(B) \geq m_i$.

In view of Properties 1–6, this can be done by distinguishing the following cases:

I: $1 \leq i \leq N$, $m_i > m_{i-1}$, $q \in (\mathcal{D}'')_{places}$, and $D_i \in (\mathcal{D}'')_{nodes}$;

II: $1 \leq i \leq N$, $m_i > m_{i-1}$, $q \in (\mathcal{D}'')_{places}$, and $D_i \in (\mathcal{D}')_{nodes}$;

III: $1 \leq i \leq N$, $m_i > m_{i-1}$, and $q \in (\mathcal{D}')_{places}$;

IV: $i = 1$ and $m_1 = m_0$;

V: $i = 0$.

For instance, assume that conditions of case I hold. Then we have

$$\begin{aligned} \mu_{\text{GE}(\hat{B})} &= \text{GE}(B) && \text{(by Remark 3(i))} \\ &\geq \text{leastInf}(q) && \text{(by Property 1)} \\ &> \text{leastInf}(D_i) && \text{(by Property 2(b))} \\ &\geq \min\{\mu < \xi \mid \mathcal{P}^*(D_i^{(\mu)}) \neq \emptyset\} && \text{(by the very definition of the leastInf of a node)} \\ &= \mu_{m_{i-1}} + 1 && \text{(by Property 3)} \\ &> \mu_{m_i-1} \end{aligned}$$

Therefore $\widehat{\text{GE}}(B) > m_i - 1$, i.e., $\widehat{\text{GE}}(B) \geq m_i$.

The remaining cases can be proved similarly, by suitably exploiting Properties 1–6.

Concerning condition (iii) of Def. 9, let $B \in (\mathcal{C})_{\text{nodes}}$ and let $q \in \mathcal{P}$ be the place such that the edge $\langle B, q \rangle$ occurs in $\mathcal{C}$. Let $\bar{\mu} < \xi$ be the ordinal such that $B = A_{\bar{\mu}}$, $q^{(\bar{\mu})} \cap \mathcal{P}^*(B^{(\bar{\mu})}) = \emptyset$ and $q^{(\bar{\mu}+1)} \cap \mathcal{P}^*(B^{(\bar{\mu})}) \neq \emptyset$. Then, since $I_{q,\lambda}^B \neq \emptyset$, we have $\bar{\mu} < \lambda$. Moreover, since $\bar{\mu} \in \text{Sal}$, we must have $\bar{\mu} = \mu_{j_0}$, for some $j_0 \in \{0, 1, \dots, \ell - 1\}$ such that $\hat{q}^{[j_0]} \cap \mathcal{P}^*(\hat{B}^{[j_0]}) = \emptyset$ and $\hat{q}^{[j_0+1]} \cap \mathcal{P}^*(\hat{B}^{[j_0]}) \neq \emptyset$. Since $\mu_{j_0} = \bar{\mu} < \lambda \leq \mu_{\hat{\lambda}}$, we have $j_0 + 1 \leq \hat{\lambda}$ and therefore $j_0 < m_0$. In particular, since $\mathcal{P}^*(B^{(\mu_{j_0})}) \neq \emptyset$, it follows that $\mathcal{P}^*(\hat{B}^{[m_0]}) \supseteq \mathcal{P}^*(\hat{B}^{[j_0]}) \neq \emptyset$, and therefore $\mathcal{P}^*(\hat{B}^{[m_0]}) \neq \emptyset$.

Since $\mathcal{P}^*(\hat{B}^{[m_j]}) \neq \emptyset$ holds by construction, for $B \in (\mathcal{D}_j)_{\text{nodes}}$ and $j = 1, \dots, N$, it follows that condition (iii) of Def. 9 is met.

Finally, by Remark 3(iii), we can assert that $\hat{\omega} = (\{\hat{q}^{[i]}\}_{q \in \mathcal{P}})_{i \leq \ell}$, with salient steps $\{\mu_i\}_{i \leq \ell}$, has a closure equipped pumping chain at $\hat{p}$. $\blacksquare$

4 A finite representation of an infinite model

In this section we show how a finite transitive partition with a related formative process can be seen as a finite representation of an infinite model.

Consider a transitive partition related to a model for a formula Φ^- of MLSSP. Assume that a formative process of such a transitive partition is equipped with a pumping chain.

Our goal consists in iterating the sequence of the pumping chain trying to preserve, roughly speaking, the satisfiability of the formula. In this way we make the resulting model invariant under expansion of the formative process, which, from a semantic point of view, means that some places can be increased without affecting the capability of the resulting transitive partition to model the formula.

Now let Φ be the previous formula enriched with new literals of the following type: $\neg \text{Finite}(x)$. Provided that at least one place for each variable x lies in the pumping chain, Φ is automatically satisfied by the transitive process which results from an infinite prolongation of the formative process obtained by repeating infinitely many times the pumping chain.

This in turn implies that the formative process with the pumping chain together with the model of Φ^- can be seen as a finite representation of an infinite model of Φ .

The following definitions serve the above strategy.

Definition 11 When Φ is a formula of MLSSPF, we denote by Φ^- the formula which results from Φ by withdrawal of all literals of the form $\neg \text{Finite}(x)$. $\square$

Definition 12 When Φ is a formula of MLSSPF, we name A COLLECTION OF INFINITE FOR Φ any subcollection Γ of $\Sigma_{\mathcal{X}_\Phi}$ which satisfies the following requirement: for any infinite clause $\neg \text{Finite}(x)$ in Φ there exists a $\sigma \in \Gamma \cap \mathfrak{S}(x)$. $\square$

These definitions enable us to summarize the content of the preceding sections into the following necessary condition for satisfiability of an MLSSPF-formula.

Theorem 3 Let Φ be a satisfiable formula of MLSSPF. Then there exist a model $\mathcal{M} \in \{\text{sets}\}^{\mathcal{X}_\Phi}$ of Φ^- and a formative process $(\Sigma_i)_{i \leq \ell}$ for $\Sigma_{\mathcal{X}_\Phi}$ such that:

- $\ell \leq f(|\mathcal{X}_\Phi|)$, where $f \in \mathbb{N}^{\mathbb{N}}$ is a specific computable function;

- the canonical board of $\mathcal{M}$ satisfies the following condition:

There exists a “collection of infinite” Γ for Φ such that to every $\sigma \in \Gamma$ there corresponds a pumping chain, closure equipped, at σ .

The main result of this section is to show that these conditions are sufficient, too. Remember that, in order to preserve the satisfiability of the MLSSP literals, we must guarantee satisfaction of the conditions 1-3 listed in Section 2.3. Hence our claim consists in detecting which structural properties two formative processes have to share in order to produce transitive partitions which model the same MLSSP literals. Then we can verify those properties with the original process and the process obtained from the previous one by repeating infinitely many times the sequence of the pumping chain. If these properties are not disrupted by prolongation of the formative process, we obtain automatically a transitive partition that simultaneously satisfies the MLSSP literals and makes the variables involved in the pumping chain infinite through one or more of their places. Thus the infinite clauses $\neg Finite(x)$ of Φ will be satisfied, because the prolongation can be performed for any pumping chain of the original formative process and the following holds

“There exists a ‘collection of infinite’ Γ for Φ such that to every $\sigma \in \Gamma$ there corresponds a pumping chain, closure equipped, at σ .”

In order to proceed with such approach, we need some technical tools which have been introduced in [Urs05]. For the reader’s convenience, we report below the essential proofs and definitions.

Definition 13 $e \in \bigcup \mathcal{P}^{(\bullet)}$ is said to be UNUSED at $\mu \leq \xi$ if $e \notin \bigcup \bigcup \mathcal{P}^{(\mu)}$, i.e., if $e \notin z$ for any $q \in \mathcal{P}$ and any $z \in q^{(\mu)}$.

We denote by $unused(q)$ the collection of all unused elements of q . □

Definition 14 An $e \in \bigcup \mathcal{P}^{(\bullet)}$ is said to be NEW at $\mu \leq \xi$ if $e \in \Delta^{(\mu)}(q)$ for some $q \in \mathcal{P}$. □

Obviously a new element is, in particular, unused.

Lemma 9 *If b is a set made of unused elements only, the same is true of $\mathcal{P}^*(\{b\} \cup A)$.*

We shall adopt the following notation. For a couple of ordinals β', β'' we denote by $[\beta', \beta'']$ the collection of ordinals $\{\beta \mid \beta' \leq \beta \leq \beta''\}$.

The following definitions are introduced to distinguish between the new sets used to imitate the old process and elements used to increase the cardinality of the places in the collection of infinity.

We say that a transitive partition Σ is equipped with a Minus-Surplus partitioning if each block q is partitioned into two sets, namely, $Surplus(q)$ and $Minus(q)$. Consistently, we can extend this notation to a formative process $(\Sigma_\mu)_{\mu \leq \xi}$. Given a node Γ , we indicate by $Minus(\Gamma^{(\mu)})$ the collection of sets

$$\{Minus(q^{(\mu)}) \mid q \in \Gamma\}.$$

Define now a Minus-Surplus partitioning for Σ_0 , and assume that for each step μ of the process a refinement of the partition $\{\Delta^{(\mu)}(q)\}_{q \in \Sigma}$ is carried out in the following way: for each $q \in \Sigma$, the set $\Delta^{(\mu)}(q)$ is split into two sets $\Delta^{(\mu)}Minus(q) \subseteq \mathcal{P}^*(Minus(A_\mu^{(\mu)}))$ and

$\Delta^{(\mu)}\text{Surplus}(q) \subseteq (\mathcal{P}^*(A_\mu^{(\mu)}) \setminus \mathcal{P}^*(\text{Minus}(A_\mu^{(\mu)})))$.
Then define inductively

$$\text{Surplus}(q^{(\mu+1)}) = \text{Surplus}(q^{(\mu)}) \cup \Delta^{(\mu)}\text{Surplus}(q)$$

and

$$\text{Minus}(q^{(\mu+1)}) = \text{Minus}(q^{(\mu)}) \cup \Delta^{(\mu)}\text{Minus}(q).$$

As for ξ , put

$$\text{Minus}(q^{(\xi)}) = \bigcup_{\mu < \xi} \text{Minus}(q^{(\mu)})$$

and, analogously,

$$\text{Surplus}(q^{(\xi)}) = \bigcup_{\mu < \xi} \text{Surplus}(q^{(\mu)})$$

If Γ is a subset of Σ , we denote by $\text{Surplus}(\Gamma)$ the set

$$\{q \in \Gamma \mid \text{Surplus}(q) \neq \emptyset\}.$$

Definition 15 Whenever a Surplus-Minus partition is defined for all blocks of a transitive partition Σ , we say that Σ is equipped with a MINUS-SURPLUS PARTITIONING, and we denote by $\text{Surplus-Minus}(\Sigma)$ the following refinement of the original partition:

$$\{\text{Minus}(q), \text{Surplus}(q) : q \in \Sigma\}.$$

It is rather obvious that $\text{Surplus-Minus}(\Sigma) \sqsubseteq \Sigma$. □

Remark 5 Easy combinatorial arguments (see [COU02, Lemma 3.1 5(b)]) show that $\mathcal{P}^*(\cdot)$ of Surplus and Minus nodes are mutually disjoint.

Whenever you have to compare two formative processes, as in dynamical systems, you have to check initial conditions of the phenomenon and the way it evolves. In our cases the initial conditions are the two starting transitive partitions and the evolutions are the couple of formative processes that begin with the above transitive partitions. Therefore the conditions of similarity are to be divided into two groups, the former on the initial conditions the latter on the evolutions.

The following requirements are to be satisfied as initial conditions of a transitive partition in order to play the role of starting point of an imitation process (as is easily seen, they are merely combinatorial).

Definition 16 Let $(\{q^{(\mu)}\}_{q \in \mathcal{P}})_{\mu \leq \xi}, (\bullet), \mathcal{T}, \mathcal{F}, \mathcal{Q}$ be a colored $\mathcal{P}$ -process, $(\widehat{\Sigma}, \widehat{\mathcal{G}})$ be a $\widehat{\Sigma}$ -board equipped with a Minus-Surplus partitioning, $q \mapsto \widehat{q}$ be a bijection from $\mathcal{P}$ to $\widehat{\mathcal{P}}$, and $\mathcal{C}$ be a closed collection of green blocks. Assume $k' < \xi$, such that (i), (vii), (viii) and (x) of Def. 17 hold in the version $\widehat{\Sigma}_{\gamma(k')} = \widehat{\Sigma}$. We say that $\widehat{\Sigma}$ WEAKLY IMITATES Σ UPWARDS, provided that the following conditions are satisfied:

(a) for all $\Gamma \subseteq \Sigma$ and $q \in \Sigma$,

$$\bigcup \text{Minus}(\widehat{\Gamma}) \in \mathcal{P}^*(\text{Minus}(\widehat{\Gamma})) \setminus \bigcup_{q \in \Sigma} \widehat{q} \quad \text{iff} \quad \bigcup \Gamma^{(k')} \in \mathcal{P}^*(\Gamma^{(k')}) \setminus \bigcup_{q \in \Sigma} q^{(k')};$$

- (b) $q \in \Gamma \wedge \text{Surplus}(q) \neq \emptyset \wedge \text{GE}(\Gamma) \geq k'$ implies $\bigcup \widehat{\Gamma} \in \mathcal{P}^*(\widehat{\Gamma}) \setminus \bigcup_{q \in \Sigma} \widehat{q}$;
 (c) if $\text{GE}(\Gamma) < k'$, then $\bigcup \Gamma^{(k')} \in q^{(k')}$ iff $\bigcup \widehat{\Gamma} \in \widehat{q}$ and $\Gamma \in \mathcal{Q}$ implies $\mathcal{P}^*(\widehat{\Gamma}) \subseteq \bigcup \widehat{\Sigma}$. $\square$

The following are instead the properties that guarantee that the processes are similar enough in order to produce transitive partitions that model the same MLSSP literals.

Definition 17 Let $(\{q^{(\mu)}\}_{q \in \mathcal{P}})_{\mu \leq \xi}, (\bullet), \mathcal{T}, \mathcal{F}, \mathcal{Q}$ be a colored $\mathcal{P}$ -process. Besides, let $(\{\widehat{q}^{[\alpha]}\}_{\widehat{q} \in \widehat{\mathcal{P}}})_{\alpha \in [\alpha', \alpha'']}$ be a simple formative processes equipped with a Minus-Surplus partitioning. Assume that $q \mapsto \widehat{q}$ is a bijection from $\mathcal{P}$ to $\widehat{\mathcal{P}}$, $\beta'' \leq \xi$, and γ is an order preserving injection from $[\beta', \beta'']$ to $[\alpha', \alpha'']$. Let $\mathcal{C}$ be a closed collection of green blocks, and $q \mapsto \widehat{q}$ be a bijection from $\mathcal{P}$ to $\widehat{\mathcal{P}}$. We say that $(\{\widehat{q}^{[\alpha]}\}_{\widehat{q} \in \widehat{\mathcal{P}}})_{\alpha \in \gamma[[\beta', \beta'']}]$ IMITATES the segment $[\beta', \beta'']$ of the process $(\{q^{(\mu)}\}_{q \in \mathcal{P}})_{\mu \leq \xi}$ if the following hold for all β in $[\beta', \beta'']$:

- (i) $|q^{(\beta)}| = |\text{Minus}^{[\gamma(\beta)]}(\widehat{q})|$;
- (ii) $|\Delta^{(\beta)}(q)| = |\Delta^{[\gamma(\beta)]}\text{Minus}(\widehat{q})|$;
- (iii) $\Delta^{[\gamma(\beta)]}\text{Surplus}(\widehat{q}) \neq \emptyset$ implies $\beta = \text{GE}(A_\beta)$, q local trash for A_β and $q \in \mathcal{C}$;
- (iv) If $\Gamma \in \mathcal{Q}$ holds, then $\mathcal{P}^*(\widehat{\Gamma}^{[\gamma(\text{GE}(\Gamma))]}]) \subseteq \bigcup \widehat{\Sigma}^{[\gamma(\text{GE}(\Gamma)+1)]}$;
- (v) For all $\beta \neq \text{GE}(\Gamma)$ $\bigcup \Gamma^{(\beta)} \in \Delta^{(\beta)}(q)$ iff $\bigcup \text{Minus} \widehat{\Gamma}^{[\gamma(\beta)]} \in \Delta^{[\gamma(\beta)]}(\widehat{q})$;
- (vi) If $\beta = \text{GE}(\Gamma)$ then $\bigcup \Gamma^{(\beta)} \in \Delta^{(\beta)}(q)$ iff $\bigcup \Gamma^{[\gamma(\beta)]} \in \Delta^{[\gamma(\beta)]}(\widehat{q})$;
- (vii) For all $q \in \mathcal{F}$ $\widehat{q}^{[\gamma(\beta)]} = \text{Minus}^{[\gamma(\beta)]}(\widehat{q})$;
- (viii) For all ordinals β $\{q : \widehat{q} \in \text{Surplus}(\widehat{\Sigma})^{[\gamma(\beta)]}\} \subseteq \mathcal{C}$;
- (ix) $|\mathcal{P}^*(\text{Minus}(\widehat{\Gamma}^{[\gamma(k)]})) \setminus \bigcup_{q \in \Sigma} q^{[k]}| = |\mathcal{P}^*(\Gamma)^{(k)} \setminus \bigcup_{q \in \Sigma} q^{(k)}|$;
- (x) $|\mathcal{P}^*(\text{Minus}(\widehat{\Gamma}^{[\gamma(k-1)]})) \cap q^{[k]}| = |\mathcal{P}^*(\Gamma^{(k-1)}) \cap q^{(k)}|$.

$\square$

Remark 6 We make some simple remarks.

- $\mathcal{P}^*(\Gamma^{(k-1)}) \cap q^{(k)} = \mathcal{P}^*(\Gamma^{(k)}) \cap q^{(k)}$. Hence, whenever $\gamma(k)$ is the successor of $\gamma(k-1)$, (x) can be rephrased as

$$|\mathcal{P}^*(\text{Minus}(\widehat{\Gamma}^{[\gamma(k)]})) \cap q^{[k]}| = |\mathcal{P}^*(\Gamma^{(k)}) \cap q^{(k)}|.$$

- Naturally, (ix) belongs to the structural properties that a formative process has to fulfill in order to simulate another one, although it can be obtained from (i) and (x).
- Assume that (viii) holds at the beginning of the process. Then (iii) entails (viii); therefore, in order to prove inductively the previous properties, it suffices to show that (viii) holds at the outset. The same argument holds for (x), which can be obtained from (ii), (iii) and (x) of the preceding step.

In a sense the initial conditions are very strong requests: in fact, whenever they are satisfied we can imitate any formative process that has one of them as starting point, as the following lemma shows.

Lemma 10 *Let $(\{q^{(\mu)}\}_{q \in \mathcal{P}})_{\mu \leq \xi}, (\bullet), \mathcal{T}, \mathcal{F}, \mathcal{Q}$ be a colored $\mathcal{P}$ -process, $(\widehat{\Sigma}, \widehat{\mathcal{G}})$ be a $\widehat{\Sigma}$ -board whose $\widehat{\Sigma}$ is equipped with a Minus-Surplus partitioning, $q \mapsto \widehat{q}$ be a bijection from $\mathcal{P}$ to $\widehat{\mathcal{P}}$, and $\mathcal{C}$ be a closed collection of green blocks. Assume that $k' \leq \xi$, and that $\widehat{\Sigma}$ weakly imitates $\Sigma_{k'}$ upwards. Define $\widehat{\Sigma} = \widehat{\Sigma}_{\gamma(k')}$ and, for all $q \in \widehat{\mathcal{P}}$, $\widehat{q} = \widehat{q}^{[\gamma(k')]}$. Then for all ordinals k'' such that $k'' \leq \xi$ and $||[k', k'']| < \omega$, one can construct a formative process $(\{\widehat{q}^{[\alpha]}\}_{\widehat{q} \in \widehat{\mathcal{P}}})_{\gamma(k') \leq \mu \leq \gamma(k'')}$ imitating the segment $[k', k'']$ of $(\{q^{(\mu)}\}_{q \in \mathcal{P}})_{\mu \leq \xi}, (\bullet), \mathcal{T}, \mathcal{F}, \mathcal{Q}$.*

Proof. We construct a formative process satisfying the conditions (i)–(x) by induction.

Concerning the base case $\mu = \gamma(k')$, (i), (vii), (viii), (x) hold by hypothesis, and (ix) holds by Remark 6, since (i) and (x) hold. Assume $k' \neq \text{GE}(A_{k'})$; then, using (ix) and hypothesis (a), we can define a partition $\bigcup_{q \in \Sigma} (\Delta^{[\gamma(k')]}(q))$ of

$$\mathcal{P}^*(\text{Minus}^{[\gamma(k')]}(\widehat{A_{k'}})) \setminus \bigcup_{q \in \Sigma} q^{[\gamma(k')]}$$

such that (ii) and (v) hold, as well. If $k' = \text{GE}(A_{k'})$ and $\text{Surplus}(\widehat{q}^{[\gamma(k')]})) \neq \emptyset$ for some $q \in A_{k'}$ (otherwise we proceed as before, and condition (vi) is automatically satisfied), then, using (b), interchanging $\bigcup \text{Minus}^{[\gamma(k')]}(\widehat{A_{k'}})$ with $\bigcup A^{[\gamma(k')]}$, (vi) is satisfied.

If $A_{k'} \in \mathcal{Q}$ and $\widehat{A_{k'}} = \text{Minus}(\widehat{A_{k'}})$, proceed as before (in this case (iv) holds by a straight checking of cardinality starting from (ix)). Otherwise, since (viii) holds, there must exist a local trash $q \in \mathcal{C}$ for $A_{k'}$. Then, construct the partition as before, except for $\Delta^{[\gamma(k')]} \text{Surplus}(\widehat{q})$, in which we put the whole remainder

$$(\mathcal{P}^*(\widehat{A_{k'}}^{[\gamma(k')]})) \setminus \bigcup_{q \in \Sigma} q^{[\gamma(k')]} \setminus \bigcup_{q \in \Sigma} \Delta^{[\gamma(k')]} \text{Minus}(\widehat{q}),$$

so satisfying (iii) and (iv).

Now, assume all the inductive hypotheses for $\gamma(k)$. Our aim is to demonstrate the case $\gamma(k+1)$. By Remark 6, provided that (iii)[$\gamma(k+1)$] is proven, (viii) automatically holds. Plainly (i)[$\gamma(k)$] and (ii)[$\gamma(k)$] entail (x)[$\gamma(k+1)$] and (i)[$\gamma(k+1)$]. The latter in its turn implies the following for all $\Gamma \subseteq \Sigma$:

$$|\mathcal{P}^*(\text{Minus}^{[\gamma(k+1)]}(\widehat{\Gamma}))| = |\mathcal{P}^*(\Gamma^{(k+1)})|. \quad (3)$$

In order to show (ix) we observe that, since

$$\begin{aligned} & \mathcal{P}^*(\text{Minus}(\widehat{\Gamma}^{[\gamma(k+1)]})) \setminus \bigcup_{q \in \Sigma} q^{[\gamma(k+1)]} \\ &= (\mathcal{P}^*(\text{Minus}(\widehat{\Gamma}^{[\gamma(k+1)]})) \setminus \mathcal{P}^*(\text{Minus}(\Gamma^{[\gamma(k)]}))) \setminus \bigcup_{q \in \Sigma} q^{[\gamma(k+1)]} \cup \\ & \cup (\mathcal{P}^*(\text{Minus}(\widehat{\Gamma}^{[\gamma(k)]})) \setminus \bigcup_{q \in \Sigma} q^{[\gamma(k+1)]}), \end{aligned}$$

it follows that

$$\begin{aligned} & \mathcal{P}^*(\text{Minus}(\widehat{\Gamma}^{[\gamma(k+1)]})) \setminus \mathcal{P}^*(\text{Minus}(\widehat{\Gamma}^{[\gamma(k)]})) \setminus \bigcup_{q \in \Sigma} q^{[\gamma(k+1)]} \\ &= \mathcal{P}^*(\text{Minus}(\widehat{\Gamma}^{[\gamma(k+1)]})) \setminus \mathcal{P}^*(\text{Minus}(\Gamma^{[\gamma(k)]})). \end{aligned}$$

Therefore,

$$\begin{aligned} & \mathcal{P}^*(\text{Minus}(\widehat{\Gamma}^{[\gamma(k+1)]})) \setminus \bigcup_{q \in \Sigma} q^{[\gamma(k+1)]} \\ &= \mathcal{P}^*(\text{Minus}(\widehat{\Gamma}^{[\gamma(k+1)]})) \setminus \mathcal{P}^*(\text{Minus}(\widehat{\Gamma}^{[\gamma(k)]})) \cup \mathcal{P}^*(\text{Minus}(\widehat{\Gamma}^{[\gamma(k)]})) \setminus \bigcup_{q \in \Sigma} q^{[\gamma(k+1)]}. \end{aligned}$$

Reasoning in the same way, we obtain

$$\begin{aligned} & \mathcal{P}^*(\Gamma)^{(k+1)} \setminus \bigcup_{q \in \Sigma} q^{(k+1)} \\ &= \mathcal{P}^*(\Gamma)^{(k+1)} \setminus \mathcal{P}^*(\Gamma)^{(k)} \cup \mathcal{P}^*(\Gamma)^{(k)} \setminus \bigcup_{q \in \Sigma} q^{(k+1)}. \end{aligned}$$

By the induction hypothesis (i)[$\gamma(k)$] we have $|\mathcal{P}^*(\text{Minus}^{[\gamma(k)]}\widehat{\Gamma})| = |\mathcal{P}^*(\Gamma^{(k)})|$ and, by equation (3),

$$|\mathcal{P}^*(\text{Minus}^{[\gamma(k+1)]}\widehat{\Gamma})| = |\mathcal{P}^*(\Gamma^{(k+1)})|,$$

which in its turn implies

$$|\mathcal{P}^*(\text{Minus}(\widehat{\Gamma}^{[\gamma(k+1)]})) \setminus \mathcal{P}^*(\text{Minus}(\widehat{\Gamma}^{[\gamma(k)]}))| = |\mathcal{P}^*(\Gamma)^{(k+1)} \setminus \mathcal{P}^*(\Gamma)^{(k)}|.$$

Hence we are left to prove the equality

$$|\mathcal{P}^*(\text{Minus}(\widehat{\Gamma}^{[\gamma(k)]})) \setminus \bigcup_{q \in \Sigma} q^{[\gamma(k+1)]}| = |\mathcal{P}^*(\Gamma)^{(k)} \setminus \bigcup_{q \in \Sigma} q^{(k+1)}|. \quad (4)$$

Observe that

$$\mathcal{P}^*(\text{Minus}(\widehat{\Gamma}^{[\gamma(k)]})) \setminus \bigcup_{q \in \Sigma} q^{[\gamma(k+1)]} = \mathcal{P}^*(\text{Minus}(\widehat{\Gamma}^{[\gamma(k)]})) \setminus \bigcup_{q \in \Sigma} q^{[\gamma(k)]} \setminus \bigcup_{q \in \Sigma} \Delta^{[\gamma(k)]}\text{Minus}(q).$$

If $\Gamma \neq A_k$, by the disjointness of $\mathcal{P}^*$ we get

$$\mathcal{P}^*(\text{Minus}(\widehat{\Gamma}^{[\gamma(k)]})) \setminus \bigcup_{q \in \Sigma} q^{[\gamma(k)]} \setminus \bigcup_{q \in \Sigma} \Delta^{[\gamma(k)]}\text{Minus}(q) = \mathcal{P}^*(\text{Minus}(\widehat{\Gamma}^{[\gamma(k)]})) \setminus \bigcup_{q \in \Sigma} q^{[\gamma(k)]}.$$

Plainly, the same is true in the $\perp^0$ version, thus (4) holds for $\gamma(k)$, by virtue of (ix). Otherwise, since $\bigcup_{q \in \Sigma} \Delta^{[\gamma(k)]}\text{Minus}(q)$ is a partition of a subset extract from

$$\mathcal{P}^*(\text{Minus}(\widehat{\Gamma}^{[\gamma(k)]})) \setminus \bigcup_{q \in \Sigma} q^{[\gamma(k)]},$$

we have that

$$\begin{aligned} & \left| \mathcal{P}^*(\text{Minus}(\widehat{\Gamma}^{[\gamma(k)]})) \setminus \bigcup_{q \in \Sigma} q^{[\gamma(k)]} \setminus \bigcup_{q \in \Sigma} \Delta^{[\gamma(k)]}\text{Minus}(q) \right| \\ &= \left| \mathcal{P}^*(\text{Minus}(\widehat{\Gamma}^{[\gamma(k)]})) \setminus \bigcup_{q \in \Sigma} q^{[\gamma(k)]} \right| - \sum_{q \in \Sigma} |\Delta^{[\gamma(k)]}\text{Minus}(q)|. \end{aligned}$$

Again, the same holds in the $\perp^0$ version, and (4) is reached by (i)[$\gamma(k)$] and (ii)[$\gamma(k)$]. This concludes the proof of (ix)[$\gamma(k+1)$].

Concerning (vii)[$\gamma(k+1)$], observe that $q^{[\gamma(k+1)]} = q^{[\gamma(k)]} \cup \Delta^{[\gamma(k)]}(q)$. By the induction hypothesis (vii)[$\gamma(k)$],

$$q^{[\gamma(k)]} = \text{Minus}(q^{[\gamma(k)]}).$$

On the other side, since (iii)[$\gamma(k)$] holds and $\mathcal{C}$ is composed of green places only,

$$\Delta^{[\gamma(k)]}(q) = \Delta^{[\gamma(k)]}\text{Minus}(q),$$

which implies (vii)[$\gamma(k+1)$].

Regarding (ii)[$\gamma(k+1)$]- (vi)[$\gamma(k+1)$], the argument goes like in the base case. $\blacksquare$

The following theorem shows that our requests are well done, that means that they are sufficient for the role we are expected to play. Indeed, they produce transitive partitions that model the same MLSSP literals.

Theorem 4 *Let $((\{q^{(\mu)}\}_{q \in \mathcal{P}})_{\mu \leq \xi}, (\bullet), \mathcal{T}, \mathcal{F}, \mathcal{Q})$ be a colored $\mathcal{P}$ -process. Moreover, let $(\{\hat{q}^{[\alpha]}\}_{\hat{q} \in \hat{\mathcal{P}}})_{\alpha \leq \xi'}$ be another formative process, equipped with a Minus-Surplus partitioning. Assume that, for some $k' \leq \xi$ and $m \leq \xi'$,*

- $\hat{\Sigma}_m$ weakly imitates $\Sigma_{k'}$ upwards;
- the process $(\{\hat{q}^{[\alpha]}\}_{\hat{q} \in \hat{\mathcal{P}}})_{\alpha \in \gamma[k', \xi]}$ imitates $(\{q^{(\mu)}\}_{q \in \mathcal{P}})_{k' \leq \mu \leq \xi}$, where γ is an injective map from $[k', \xi]$ to $[m, \xi']$;
- $\hat{\Sigma}_{\xi'}$ has the same targets as Σ_{ξ} ;
- for all $\mu > m$ such that $\mu \notin \gamma[k', \xi]$, the following holds: $\Delta^{[\mu]}(\hat{q}) \subseteq \Delta^{[\mu]}\text{Surplus}(\hat{q})$;
- if β is the largest ordinal such that $\beta \in \gamma[k', \xi] \wedge \beta \leq \mu$, if q is a local trash of A_μ , and if $\text{GE}(A_\mu) > \gamma^{-1}(\beta)$, then $\bigcup \widehat{A}_\mu^{[\xi']} \notin \Delta^{[\mu]}\text{Surplus}(\hat{q})$.

Consider an MLSPF-formula Φ , a set-valued assignment $\mathcal{M} \in \{\text{sets}\}^{\mathcal{X}_\Phi}$ defined on the collection $\mathcal{X}_\Phi$ of variables in Φ , and assume that $((\{q^{(\mu)}\}_{q \in \mathcal{P}})_{\mu \leq \xi}, (\bullet), \mathcal{T}, \mathcal{F}, \mathcal{Q})$ is a colored $\mathcal{P}$ -process for the $\Sigma_{\mathcal{X}_\Phi}$ -board.

Then, letting $\mathcal{M}'(v) = \bigcup [\mathfrak{S}_{\mathcal{M}}(\hat{v})]$ for every literal in Φ , the following conditions are fulfilled:

- if the literal is satisfied by $\mathcal{M}$, then it is satisfied by $\mathcal{M}'$ too;
- if the literal is satisfied by $\mathcal{M}'$, and does not involve $\mathcal{P}$ or the construct $\{-, \dots, -\}$, then it is satisfied by $\mathcal{M}$ too.

The proof of Theorem 4 can be found in [Urs05] but, for the sake of completeness, we review it in the Appendix.

Now we are ready for the main result of this section. We show how a finite assignment can yield an infinite model, which satisfies the same MLSSP literals. To this end, we prove the following theorem, which generalizes a result given in [Urs05, Theorem 38].

Theorem 5 *Let Φ be a formula of MLSPF and $\mathcal{M} \in \{\text{sets}\}^{\mathcal{X}_\Phi}$ a model for Φ^- together with a $((\{q^{(\mu)}\}_{q \in \mathcal{P}})_{\mu \leq \ell}, (\bullet), \mathcal{T}, \mathcal{F}, \mathcal{Q})$ colored $\mathcal{P}$ -process for $\Sigma_{\mathcal{X}_\Phi}$ such that:*

- $\mathcal{G} = (\mathcal{T}, \mathcal{F}, \mathcal{Q})$ colored $\Sigma_{\mathcal{X}_\Phi}$ -board satisfies the following conditions:

There exist a “collection of infinite” Γ for Φ such that for all $\sigma \in \Gamma$ there exist a pumping chain, closure equipped, at σ . $\langle (\mathcal{C}, \overline{\mathcal{D}}_0, \mathcal{D}_1, \dots, \mathcal{D}_N), \vec{m}, q_0, \sigma \rangle$ with a closure $\overline{\mathcal{C}}$.

then there exists a model for Φ .

Proof. Consider an occurrence of the literal $\neg Finite(x)$.

Since Γ is a collection of infinite there exist a place $\sigma \subseteq \mathcal{M}(x)$ and a pumping chain, $\langle (\mathcal{C}, \overline{\mathcal{D}}_0, \mathcal{D}_1, \dots, \mathcal{D}_N), \vec{m}, q_0, \sigma \rangle$ with a closure $\overline{\mathcal{C}}$.

Our goal consists in building from the original $((\{q^{(\mu)}\}_{q \in \mathcal{P}})_{\mu \leq \ell}, (\bullet), \mathcal{T}, \mathcal{F}, \mathcal{Q})$ colored $\mathcal{P}$ -process a new one $(\{\hat{q}^{[\alpha]}\}_{\hat{q} \in \hat{\mathcal{P}}})_{\alpha \leq \xi'}$ in such a way $\widehat{\mathcal{M}}(v) = \bigcup \{\hat{q} : q \in \mathcal{M}(v)\}$ is a model for Φ .

We define, using a technique described in [CU97], a sequence of nodes of length ξ , denoting it $\mathcal{A}_\xi$. We shall use such a sequence as trace for a formative process with final transitive partition $\widehat{\mathcal{P}}$ satisfying the requested properties. The following sequence does to the purpose:

$$A_1 \dots A_{m_0-1} \underbrace{\overbrace{C_0 \dots C_t}^{\mathcal{C}} \overbrace{D_0 \dots D_{s_1-1}}^{\overline{\mathcal{D}}_0} \overbrace{C_{t+1} \dots C_h}^{\mathcal{C}}}_{\text{N}_0\text{-times}} A_{\gamma(m_0)} \dots A_{\gamma(m_1-1)} \\ \underbrace{D_{s_0=0} \dots D_{s_1-1}}^{\mathcal{D}_1} A_{\gamma(m_1)} \dots A_{\gamma(m_N-1)} \underbrace{D_{s_{N-1}} \dots D_{s_N}}^{\mathcal{D}_N} A_{\gamma(m_N)} \dots A_{\gamma(\xi)}.$$

Concerning nodes of the original sequence we use the capital letter A , concerning nodes of the cycle $\mathcal{C}$ we use the capital letter C , and concerning nodes of the simple path $\mathcal{D}$ attached to $\mathcal{C}$ we use the capital letter D . We denote by:

- N_1 , the nodes of the sequence $\mathcal{A}_l$;
- N_2 , the nodes of the pumping cycle $(\mathcal{C}, \mathcal{D})$;
- $A_{\gamma(i)}$, a node A_i in the pumped sequence;
- $A_{\gamma(i)+t}$, an N_2 -node, where $\gamma(i)$ is the index of the first N_1 -node before $A_{\gamma(i)+t}$.

Finally, we usually refer to a piece of a formative process, bounded by the two indices $\alpha \leq \beta$, by “segment of the process”. We refer to the segments composed of N_1 and N_2 nodes in a different way, the former as “imitating segments”, the latter as “pumping segments”.

In order to define a formative process we are needed only to exhibit the way to distribute all the elements produced at each stage. Our strategy consists in copying the old formative process until the stage $m_0 - 1 = \gamma(m_0 - 1)$, therefore along this segment we define γ as the Identity map; then “pumping” the cycle to create new elements and distribute them. This procedure by transfinite induction increases the cardinality of the blocks inside the cycle. Unfortunately this distribution of elements could be not propagated along the path $\mathcal{D}$ in order to make σ of infinite size. This phenomenon depends on the fact that some of the nodes involved could be not “ready” at the step when one of its block is infinite. This obliges us to wait for the first step when it is possible. As soon as these conditions are reached the “infinite can be propagated” until making σ of infinite size. This procedure has in mind to preserve the cardinality of all blocks not involved in the pumping procedure. In order to do this we distinguish the elements reserved for pumping procedure from those used for mimicking the old process, creating a Surplus-Minus partition inside all the blocks involved in the pumping procedure, actually refining the original partition.

Assume we have copied the original formative process until $m_0 - 1$.

We start with the stage $\gamma(m_0 - 1) + 1$.

We trigger off our pumping procedure using the property (i) of the definition of pumping chain. Indeed, by (i) $\mathcal{C}$ has an q_0 block with a new element t_0 .

This element, t_0 , will generate all the elements needed for increasing the size of all the blocks inside the cycle.

[For the details of such a construction, cf. the proof of Theorem 38 in [Urs05]].

From now on we denote by q places in $\mathcal{C}$, by δ places in $(\overline{\mathcal{D}}_0)_{places}$, by σ_Γ a place which lies in the pumping path after the node Γ , by σ^Γ a place which lies in the pumping path before the node Γ . Put $\mathcal{D}_0 \equiv_{\text{Def}} \mathcal{C}, \overline{\mathcal{D}}_0$. It is possible to prolongate the formative process by repeating the nodes in $(\mathcal{C}, \overline{\mathcal{D}}_0)_{nodes}$ $\aleph_0$ -times in the following sequence:

$$C_0 \dots C_t, D_0 \dots D_{s_1-1}, C_{t+1} \dots C_h$$

in such a way as to enforce that the following holds at step $\gamma(m_0)$:

- (i) For all $\sigma \in (\mathcal{D}_0)_{places}$ and $\Gamma \in (\mathcal{D}_0)_{nodes}$,
 $|\mathcal{P}^*(\{\text{Surplus}^{[\gamma(m_0)]}(\sigma^\Gamma)\} \cup (\Gamma \setminus \{\sigma^\Gamma\}) \cap \text{Surplus}^{[\gamma(m_0)]} q_\Gamma| \geq \omega$;
- (ii) for all q , $\text{Minus}^{[\gamma(m_0-1)]} q = \text{Minus}^{[\gamma(m_0)]} q =_c q^{(m_0)}$ and
 $|\text{Unused}(\text{Minus}^{[\gamma(m_0)]} q)| = |\text{Unused}(q)|$;
- (iii) for all q , $\text{Surplus}(q) \neq \emptyset$ implies $q \in \text{Closure}((\mathcal{C}, \mathcal{D}))$;
- (iv) for all Γ in $(\mathcal{C}, \mathcal{D})_{nodes}$, $\bigcup \Gamma^{[\gamma(m_0)]} \notin \mathcal{P}^{[\gamma(m_0)]}$.

It is straightforward to verify that the conditions requested in the Lemma 10 are respectively satisfied by the *colored $\mathcal{P}$ -process* $(\Sigma_\mu)_{\mu \leq \xi}, (\bullet), \mathcal{T}, \mathcal{F}, \mathcal{Q}$ and the partition $\mathcal{P}^{[\gamma(m_0)]}$ for $k = m_0$. Therefore the segment $[m_0, m_1 - 1]$ can be imitated.

Using the fact that partitions Surplus are never used along the segment $[\gamma(m_0), \gamma(m_1 - 1)]$ and that it imitates the segment $[m_0, m_1 - 1]$, we can assert that for Γ in $(\mathcal{D})_{nodes}$ $\mathcal{P}^*(\Gamma)^{[\gamma(m_1)-1]} \neq \emptyset$ and $|\text{Surplus}^{[\gamma(m_1)-1]} \delta_{s_1}| \geq \omega$. This enables us to prolongate the process in such a way that an analogue of (i)–(iv) holds at the step $\gamma(m_1)$ (with $\gamma(m_1)$ in the segment $\mathcal{D}_1$). Lemma 10 applies again, and the process can be prolonged till $\gamma(m_2 - 1)$.

In the same fashion the construction proceeds by application the same procedure to every section of the pumping chain, until the last node of the trace $\mathcal{A}_\xi$ is reached. This plainly gives $|\widehat{\sigma}^{[\xi]}| \geq \omega$.

Now we are left to prove that the resulting transitive partition $\widehat{\mathcal{P}}^{[\xi]}$ satisfies the thesis.

To do this we invoke Theorem 4. It is routine verifying that the hypotheses of Theorem 4 are met. Therefore we can assert that the model induced by the resulting transitive partition satisfies Φ^- plus all the literals $\neg \text{Finite}(x)$ such that in the original model $\mathcal{M}(x) \supseteq \sigma$ are satisfied. Indeed, the previous procedure has made σ of infinite size. We are left to do the same for the other members of the collection of infinite. This is hardly ever a limitation since all pumping chains and closures are preserved and the formative process $\widehat{\Sigma}$ can do the same role played by the original process before. Therefore, the same technique applies using a new pumping chain, till the end.

Denote the last model $\mathcal{M}$ and put $X = \{x \mid \sigma \in \Gamma \cap \mathcal{M}(x)\}$.

$\mathcal{M}$ will satisfy the following formula

$$\Phi' = \Phi^- \wedge \bigwedge_{x \in X} \neg \text{Finite}(x)$$

since Γ is a collection of infinite the thesis plainly follows. ■

Combining the results of the two sections, we achieve the main conclusion of this paper:

Theorem 6 *The language MLSSPF, namely MLSSP extended with a finiteness predicate, has a decidable satisfiability problem.*

5 Conclusions

We have shown that the extension MLSSPF of the language MLSSP with a finiteness predicate has a decidable satisfiability problem, even though it does not enjoy the small model property, as is the case with MLSSP. Such result has been achieved by showing that MLSSPF enjoys a weaker form of the finite model property, called “small model witness property.” Roughly speaking, the small model witness property for MLSSPF asserts that any satisfiable MLSSPF-formula has a model admitting a “small” representation through a suitably annotated formative process.

We plan to further extend the techniques introduced in this paper to deal also with set-theoretic fragments involving besides the Boolean set operators and the singleton operator, also the unionset or the Cartesian product operators. Notice that in such fragments it is possible to express formulae admitting only infinite models.

Appendix

A Proof of Theorem 4

The introduction of some additional tools makes the proof of the Theorem 4 more plain and fluent. We create various levels of similarity of transitive partitions and formative processes and break the proof in more steps. In the following we use conditions 1,2 and 4, defined in Section 2.3, in order to define a new class of similarity among transitive partitions.

Definition 18 Let $\mathcal{G}$ be a colored transitive Σ -board. Then $\widehat{\Sigma}$ is said to **SIMULATE** $(\Sigma, \mathcal{G})$ **UPWARDS**, when there is a bijection $\beta \in \widehat{\Sigma}^\Sigma$ such that

- $\widehat{\Sigma}$ $\in$ -simulates Σ via β . That is, $\bigcup \beta[X] \in \bigcup \beta[Y]$ if and only if $\bigcup X \in \bigcup Y$, for $X, Y \subseteq \Sigma$;
- $\widehat{\Sigma}$ $\mathcal{P}$ -simulates Σ via β . That is, $\bigcup \beta[X] = \mathcal{P}(\bigcup \beta[Y])$ if $\bigcup X = \mathcal{P}(\bigcup Y)$, for $Y \in \mathcal{Q}$ $X, Y \subseteq \Sigma$;
- $\widehat{\Sigma}$ *Red*-simulates Σ via β . That is, if $\sigma \in \mathcal{F}$, then $|\beta(\sigma)| = |\sigma|$.

□

The following lemma shows that the previous conditions are enough to model **MLSSP** clauses.

Lemma 11 Consider a formula $\Phi \in \text{MLSSPF}$, a set-valued assignment $\mathcal{M} \in \{\text{sets}\}^{\mathcal{X}_\Phi}$ defined on the collection $\mathcal{X}_\Phi$ of variables in Φ , together with the colored transitive $\Sigma_{\mathcal{X}_\Phi}$ -board $\mathcal{G} = (\mathcal{T}, \mathcal{F}, \mathcal{Q})$. Define Φ^- as the formula Φ without literals of the type *Finite*(x) or $\neg \text{Finite}(x)$. Moreover, let $\widehat{\Sigma}$ be a partition and β be a bijection between $\Sigma_{\mathcal{X}_\Phi}$ and $\widehat{\Sigma}$ such that $\widehat{\Sigma}$ simulates $(\Sigma, \mathcal{G})$ upwards via β , and let $\mathcal{M}'(v) = \bigcup \beta[\mathfrak{S}_{\mathcal{M}}(v)]$. Then, for every literal in Φ^- , the following conditions are fulfilled:

- if the literal is satisfied by $\mathcal{M}$, then it is satisfied by $\mathcal{M}'$ too;
- if the literal is satisfied by $\mathcal{M}'$, and does not involve $\mathcal{P}$ or the construct $\{_, \dots, _ \}$, then it is satisfied by $\mathcal{M}$ too;
- if the literal *Finite*(x) appears in Φ and is satisfied by $\mathcal{M}$, then it is satisfied by $\mathcal{M}'$ too.

Proof. The thesis can be recast as follows. For u, v, w and w_i in $\mathcal{X}_\Phi$, the following conditions hold for all literals in Φ :

- (1) $\bigcup \mathfrak{S}(v) \mathfrak{R} \bigcup \mathfrak{S}(w)$ iff $\bigcup \beta[\mathfrak{S}(v)] \mathfrak{R} \bigcup \beta[\mathfrak{S}(w)]$, for $\mathfrak{R}$ in $\{=, \in, \subseteq\}$;
- (2) $\bigcup \mathfrak{S}(v) = \bigcup \mathfrak{S}(u) \star \bigcup \mathfrak{S}(w)$ iff $\bigcup \beta[\mathfrak{S}(v)] = \bigcup \beta[\mathfrak{S}(u)] \star \bigcup \beta[\mathfrak{S}(w)]$, for $\star$ in $\{\cap, \setminus, \cup\}$, and $\bigcup \mathfrak{S}(v) = \emptyset$ iff $\bigcup \beta[\mathfrak{S}(v)] = \emptyset$;
- (3) if $\bigcup \mathfrak{S}(v) = \mathcal{P}(\bigcup \mathfrak{S}(w))$, then $\bigcup \beta[\mathfrak{S}(v)] = \mathcal{P}(\bigcup \beta[\mathfrak{S}(w)])$;
- (4) if $\bigcup \mathfrak{S}(v) = \{\bigcup \mathfrak{S}(w_1), \dots, \bigcup \mathfrak{S}(w_H)\}$, then $\bigcup \beta[\mathfrak{S}(v)] = \{\bigcup \beta[\mathfrak{S}(w_1)], \dots, \bigcup \beta[\mathfrak{S}(w_H)]\}$;
- (5) if *Finite*(v) appears in Φ then $|\bigcup \mathfrak{S}(v)| = |\bigcup \beta[\mathfrak{S}(v)]|$.

Property (1)_∈ (here $\mathfrak{R}$ is meant to be $\in$) follows from $\in$ -simulates in Def. 18. (3) follows from the assumption $\mathfrak{S}(v) \in \mathcal{Q}$ and the notion of $\mathcal{P}$ -simulates given in the same definition. Condition (5) plainly follows from definition of *Red*-simulates.

We are left to prove that (4) hold. Observe that $\mathfrak{S}(v) \subseteq \mathcal{F}$, then consider $\mathfrak{S}(v)$ as the set X and Y_i as the sets $\mathfrak{S}(w_i)$. Hence we can assume that $\bigcup X = \{\bigcup Y_1, \dots, \bigcup Y_L\}$, $X \subseteq \mathcal{F}$, and $Y_1, \dots, Y_L$ are distinct. We must check that $\bigcup \beta[X] = \{\bigcup \beta[Y_1], \dots, \bigcup \beta[Y_L]\}$. Since $\widehat{\Sigma}$ *Red*-simulates $(\Sigma, \mathcal{G})$ and $X \subseteq \mathcal{F}$, and $|\beta(\sigma)| = |\sigma|$ for each $\sigma \in X$, the desired conclusion easily follows. Indeed, by property (1) of Def. 18, $\bigcup \beta[Y_i] \in \beta(\sigma)$ if and only if $\bigcup Y_i \in \sigma$, and $\beta[Y_1], \dots, \beta[Y_L]$ (and, accordingly, $\bigcup \beta[Y_1], \dots, \bigcup \beta[Y_L]$) are pairwise distinct.

The proofs of remaining bi-implications go exactly as in [COU02, Lemma 10.1] $\blacksquare$

The next definition is a level of similarity of transitive partitions defined in a more combinatorial way; in the sequel we show how these definitions are interrelated.

Definition 19 Consider a colored Σ -board $\mathcal{G} = (\mathcal{T}, \mathcal{F}, \mathcal{Q})$. A partition $\widehat{\Sigma}$ is said to *IMITATE* $(\Sigma, \mathcal{G})$ when there is a bijection $\beta \in \widehat{\Sigma}^\Sigma$ such that, for $\Gamma \subseteq \Sigma$, $\sigma \in \Sigma$,

- (1) $\beta(\sigma) \cap \mathcal{P}^*(\beta[\Gamma]) \neq \emptyset$ holds [if and] only if $\sigma \cap \mathcal{P}^*(\Gamma) \neq \emptyset$;
- (2) $\bigcup \beta[\Gamma] \in \beta(\sigma)$ holds if and only if $\bigcup \Gamma \in \sigma$;
- (3) if $\Gamma \in \mathcal{Q}$ holds, then $\mathcal{P}^*(\beta[\Gamma]) \subseteq \bigcup \widehat{\Sigma}$;
- (4) if $\sigma \in \mathcal{F}$ holds, then $|\beta(\sigma)| < \aleph_0$.

We will say that $\widehat{\Sigma}$ *IMITATES* $(\Sigma, \mathcal{G})$ *UPWARDS* when the following additional condition holds, for all $\sigma \in \Sigma$:

- (4') if $\sigma \in \mathcal{F}$, then $|\beta(\sigma)| = |\sigma|$.

$\square$

Lemma 12 Consider a colored Σ -board $\mathcal{G} = (\mathcal{T}, \mathcal{F}, \mathcal{Q})$ assume that a transitive partition $\widehat{\Sigma}$ *imitates* $(\Sigma, \mathcal{G})$ *upwards* then it *simulates* $(\Sigma, \mathcal{G})$ *upwards*.

Proof. Let Σ and $\widehat{\Sigma}$ be transitive partitions, and let $\mathcal{G}$ be a colored $\mathcal{P}$ -board induced by Σ . Assume that $\widehat{\Sigma}$ *imitates* $(\Sigma, \mathcal{G})$ *upwards* via the bijection $\beta \in (\widehat{\Sigma})^\Sigma$. Finally, let $X, Y \subseteq \Sigma$.

Then we have: $\bigcup \beta[X] \in \bigcup \beta[Y]$ iff $(\exists \widehat{\sigma} \in \beta[Y])(\bigcup \beta[X] \in \widehat{\sigma})$ iff $(\exists \sigma \in Y)(\bigcup \beta[X] \in \beta(\sigma))$ iff $(\exists \sigma \in Y)(\bigcup X \in \sigma)$ iff $\bigcup X \in \bigcup Y$.

Assuming now that $\bigcup X = \mathcal{P}(\bigcup Y)$, $Y \in \mathcal{Q}$, let us prove that $\mathcal{P}(\bigcup \beta[Y]) \subseteq \bigcup \beta[X]$. Indeed, suppose $t \subseteq \bigcup \beta[Y]$ and let $\widehat{\Sigma}_t$ be the subset of $\widehat{\Sigma}$ for which $t \in \mathcal{P}^*(\widehat{\Sigma}_t)$ (so that $\widehat{\Sigma}_t \subseteq \beta[Y]$, which implies $\widehat{\Sigma}_t \in \mathcal{Q}$ by the hereditarily closedness by inclusion of $\mathcal{Q}$). As $\beta^{-1}[\widehat{\Sigma}_t] \subseteq Y$, it follows that $\mathcal{P}^*(\beta^{-1}[\widehat{\Sigma}_t]) \subseteq \mathcal{P}(\bigcup Y) = \bigcup X \subseteq \bigcup \Sigma$. Therefore, by the fact that $\widehat{\Sigma}$ *imitates* $(\Sigma, \mathcal{G})$ *upwards* and $\widehat{\Sigma}_t \in \mathcal{Q}$, it follows that $\mathcal{P}^*(\widehat{\Sigma}_t) \subseteq \bigcup \widehat{\Sigma}$, so that $t \in \bigcup \widehat{\Sigma}$. Let $\widehat{\sigma}_t$ be the block in $\widehat{\Sigma}$ to which t belongs, and let σ_t be the block in Σ for which $\beta(\sigma_t) = \widehat{\sigma}_t$. Then, since $\mathcal{P}^*(\widehat{\Sigma}_t) \cap \widehat{\sigma}_t \neq \emptyset$, we have that $\mathcal{P}^*(\beta^{-1}[\widehat{\Sigma}_t]) \cap \sigma_t \neq \emptyset$, which yields $\bigcup X = \mathcal{P}(\bigcup Y) \supseteq \mathcal{P}^*(\beta^{-1}[\widehat{\Sigma}_t])$, $\bigcup X \cap \sigma_t \neq \emptyset$, $\sigma_t \in X$, and hence $t \in \widehat{\sigma}_t \in \beta[X]$, which in turn yields $t \in \bigcup \beta[X]$.

Next, assuming again $\bigcup X = \mathcal{P}(\bigcup Y)$, let us prove that $\bigcup \beta[X] \subseteq \mathcal{P}(\bigcup \beta[Y])$. Indeed, for each $t \in \bigcup \beta[X]$ there is a unique $\sigma_t \in X$ such that $t \in \beta(\sigma_t)$; moreover, by the transitivity of $\bigcup \widehat{\Sigma}$, there is a unique $\Gamma \subseteq \Sigma$ for which $t \in \mathcal{P}^*(\beta[\Gamma])$. Moreover, since $\mathcal{P}^*(\beta[\Gamma]) \cap \beta(\sigma_t) \neq \emptyset$, we also have that $\mathcal{P}^*(\Gamma) \cap \sigma_t \neq \emptyset$. Thus we can take $t' \in \sigma_t \cap \mathcal{P}^*(\Gamma)$

that, as $\sigma_t \subseteq \bigcup X = \mathcal{P}(\bigcup Y)$, fulfills $t' \in \mathcal{P}^*(Z)$ for a suitable $Z \subseteq Y$. In conclusion, $\Gamma = Z$, and therefore $t \subseteq \bigcup \beta[\Gamma] = \bigcup \beta[Z] \subseteq \bigcup \beta[Y]$. ■

As an immediate consequence, we have

Corollary 1 *Consider a formula $\Phi \in \text{MLSSPF}$, a set-valued assignment $\mathcal{M} \in \{\text{sets}\}^{\mathcal{X}_\Phi}$ defined on the collection $\mathcal{X}_\Phi$ of variables in Φ , together with the colored transitive $\Sigma_{\mathcal{X}_\Phi}$ -board $\mathcal{G} = (\mathcal{T}, \mathcal{F}, \mathcal{Q})$. Moreover, let $\widehat{\Sigma}$ and β be a partition and a bijection, respectively, such that $\widehat{\Sigma}$ imitates $(\Sigma, \mathcal{G})$ upwards via β , and let $\mathcal{M}'(v) = \bigcup \beta[\mathfrak{S}_{\mathcal{M}}(v)]$, where $\mathfrak{S}$ is the function $\mathfrak{S} \in \mathcal{P}(\Sigma)^{\mathcal{X}_\Phi}$ such that $\mathcal{M}(v) = \bigcup \mathfrak{S}(v)$ holds for every v in $\mathcal{X}$. Then, for every literal in Φ^- and any literal of the form $\text{Finite}(x)$, the following conditions are fulfilled:*

- *if the literal is satisfied by $\mathcal{M}$, then it is satisfied by $\mathcal{M}'$ too;*
- *if the literal is satisfied by $\mathcal{M}'$, and does not involve $\mathcal{P}$ or the construct $\{-, \dots, _ \}$, then it is satisfied by $\mathcal{M}$ too.* ■

We are almost ready for the desired proof of Theorem 4. One more lemma is needed showing that, provided that the initial conditions (relative to the starting transitive partition) are “similar” in the sense described above, the process which imitates the initial model satisfies the same MLSSP clauses.

Lemma 13 *Let $(\{q^{(\mu)}\}_{q \in \mathcal{P}})_{\mu \leq \xi}, (\bullet), \mathcal{T}, \mathcal{F}, \mathcal{Q}$ be a colored $\mathcal{P}$ -process. Moreover, let $(\{\widehat{q}^{[\alpha]}\}_{\widehat{q} \in \widehat{\mathcal{P}}})_{\alpha \leq \xi'}$ be another formative process, equipped with a Minus-Surplus partitioning. Assume that, for some $k' \leq \xi$ and $m \leq \xi'$,*

- $\widehat{\Sigma}_m$ weakly imitates $\Sigma_{k'}$ upwards;
- the process $(\{\widehat{q}^{[\alpha]}\}_{\widehat{q} \in \widehat{\mathcal{P}}})_{\alpha \in \gamma[k', \xi]}$ imitates $(\{q^{(\mu)}\}_{q \in \mathcal{P}})_{k' \leq \mu \leq \xi}$, where γ is an injective map from $[k', \xi]$ to $[m, \xi']$;
- $\widehat{\Sigma}_{\xi'}$ has the same targets of Σ_ξ ;
- for all $\mu > m \wedge \mu \notin \gamma[k', \xi]$ the following holds: $\Delta^{[\mu]}(\widehat{q}) \subseteq \Delta^{[\mu]} \text{Surplus}(\widehat{q})$;
- if β is the greatest ordinal such that $\beta \in \gamma[k', \xi] \wedge \beta \leq \mu$, if q is a local trash of A_μ , and if $\text{GE}(A_\mu) > \gamma^{-1}(\beta)$, then $\bigcup \widehat{A}_\mu^{[\xi']} \notin \Delta^{[\mu]} \text{Surplus}(\widehat{q})$.

Then $\widehat{\Sigma}_{\xi'}$ imitates Σ_ξ upwards.

Proof. We prove that the resulting partition $\widehat{\Sigma}_{\xi'}$ fulfills the conditions:

- (0) $q^{(\xi)} \cap \mathcal{P}^*(\Gamma)^{(\xi)} \neq \emptyset$ holds if and only if $\widehat{q}^{[\xi']} \cap \mathcal{P}^*(\widehat{\Gamma}^{[\xi']}) \neq \emptyset$;
- (1) $\bigcup \widehat{\Gamma}^{[\xi']} \in \widehat{q}^{[\xi']}$ if and only if $\bigcup \Gamma^{(\xi)} \in q^{(\xi)}$;
- (2) if $\Gamma \in \mathcal{Q}$ holds, then $\mathcal{P}^*(\widehat{\Gamma}^{[\xi']}) \subseteq \bigcup \widehat{\mathcal{P}}^{[\xi']}$;
- (3') if $q \in \mathcal{F}$, then $|\widehat{q}^{[\xi']}| = |q^{(\xi)}|$.

Along the verification of properties (0)-(3') we refer to (i)-(x) of Def. 17.

- (0) By the fact that the two partitions have the same targets;

- (1) In case $\bigcup \widehat{\Gamma}^{[\xi']} \in \widehat{q}^{[\xi']}$, assuming that it is distributed strictly before m , then $\text{GE}(\Gamma) < k'$. Indeed, if not so, by (vi) Def. 17, since $\bigcup \Gamma^{(\text{GE}(\Gamma))} \in \Delta^{(\text{GE}(\Gamma))}(q)$,

$$\bigcup \widehat{\Gamma}^{[\xi']} \in \widehat{q}^{[\xi']} = \bigcup \widehat{\Gamma}^{[\gamma(\text{GE}(\Gamma))]} \in \Delta^{[\gamma(\text{GE}(\Gamma))]} \widehat{q},$$

which is impossible, due to the fact that $\bigcup \widehat{\Gamma}^{[\xi']} \in \widehat{q}^{[\xi']}$ is already in $\widehat{q}^{[\gamma(\text{GE}(\Gamma))]}$, and $\Delta^{[\gamma(\text{GE}(\Gamma))]} \widehat{q}$, by definition, is made of elements of $\mathcal{P}^*(\widehat{\Gamma}^{[\gamma(\text{GE}(\Gamma))]} \setminus \bigcup_{q \in \Sigma} q^{[\gamma(\text{GE}(\Gamma))]}).$

Then, using the fact that $\widehat{\Sigma}_m$ weakly simulates $\Sigma_{k'}$, the result follows. Concerning the right implication, we are left to prove the case when $\bigcup \widehat{\Gamma}^{[\xi']}$ is distributed after or in m . Let j be such an index. By hypothesis, j cannot be outside $\gamma[k', \xi]$, and so $j = \gamma(k)$ for some k . We show that $k = \text{GE}(\Gamma)$. By contradiction, let us assume $k > \text{GE}(\Gamma)$. Then, by (vi) Def. 17,

$$\bigcup \widehat{\Gamma}^{[\gamma(\text{GE}(\Gamma))]} \in \Delta^{[\gamma(\text{GE}(\Gamma))]} \widehat{q}.$$

Observe that, after $\gamma(\text{GE}(\Gamma))$, $\widehat{\Gamma}$ cannot change inside the range of γ , on account of (ii) and (iii) of Def. 17. It cannot change for an index j outside, since $\text{GE}(\Gamma)$ is greater than the greatest ordinal β such that $\beta \in \gamma[k', \xi] \wedge \beta \leq j$. On the other hand, k cannot be strictly less than $\text{GE}(\Gamma)$, since in this case the same argument used for $\bigcup \widehat{\Gamma}^{[\xi']}$ distributed before m and $\text{GE}(\Gamma) \geq k'$ applies. Therefore $k = \text{GE}(\Gamma)$, and we are done. We now show the left implication in the case $\text{GE}(\Gamma) < k'$. The hypothesis implies that $\bigcup \widehat{\Gamma}^{[m]} \in \widehat{q}^{[m]}$. Reasoning as before, we conclude that $\widehat{\Gamma}$ cannot change along the process after m . Finally, assuming $\text{GE}(\Gamma) \geq k'$, by Def. 17(vi) we have

$$\bigcup \widehat{\Gamma}^{[\gamma(\text{GE}(\Gamma))]} \in \Delta^{[\gamma(\text{GE}(\Gamma))]} \widehat{q}.$$

Again $\widehat{\Gamma}$ cannot change in the sequel of the process, either along the imitated process, or outside.

- (2) Follows plainly from Def. 17(iv). Indeed, $\Gamma \in \mathcal{Q}$, therefore

$$\mathcal{P}^*(\widehat{\Gamma}^{[\gamma(\text{GE}(\Gamma))]} \subseteq \bigcup \widehat{\mathcal{P}}^{[\gamma(\text{GE}(\Gamma))]}.$$

As observed in the previous point, after $[\gamma(\text{GE}(\Gamma))]$, $\widehat{\Gamma}$ cannot change either along the imitating process, by (ii) and Def. 17(iii), or outside, by hypothesis. Thus $\mathcal{P}^*(\widehat{\Gamma}^{[\xi']}) \subseteq \bigcup \widehat{\mathcal{P}}^{[\xi']}$.

- (3') The red places cannot belong to $\mathcal{C}$. Hence, by the property (viii), they cannot have Surplus part, which in turns implies that $\text{Minus}(\widehat{q}^{[\xi']}) = \widehat{q}^{[\xi']}$. This, combined with $|\text{Minus}^{[\gamma(\xi)]}(q)| = |q^{(\xi)}|$, by Def. 17(i), yields the thesis. ■

By combining the results in Lemma 13 and Corollary 1, Theorem 4 follows easily.

References

- [Can91] D. Cantone, Decision procedures for elementary sublanguages of Set Theory: X. Multi-level syllogistic extended by the singleton and powerset operators. *J. Automat. Reason.* **7**(2):193–230, 1991.
- [CFO89] D. Cantone, A. Ferro, and E.G. Omodeo. *Computable Set Theory*. Vol. 6 Oxford Science Publications of *International Series of Monographs in Computer Science*. Clarendon Press, Oxford, 1989.
- [COP01] D. Cantone, E. G. Omodeo, and A. Policriti. *Set Theory for Computing — From decision procedures to declarative programming with sets*. Monographs in Computer Science. Springer-Verlag, New York, 2001.
- [COS*03] D. Cantone, E. G. Omodeo, J. T. Schwartz, and P. Ursino. Notes from the logbook of a proof-checker’s project. In N. Dershowitz, editor, *Proc. of the International Symposium on Verification: Theory and Practice*, volume 2772 of *Lecture Notes in Computer Science*, pages 182–207, Berlin, Springer-Verlag, 2003.
- [COU02] D. Cantone, E. G. Omodeo, and P. Ursino. Formative processes with applications to the decision problem in set theory: I. Powerset and singleton operators. *Inform. and Comput.*, **172**:165–201, 2002.
- [CU97] D. Cantone and P. Ursino. A unifying approach to computable set theory. *Proceedings of Logic Colloquium 1997*, Leeds, UK, 6 - 13 July 1997.
- [Jec78] T. Jech. *Set theory*, Academic Press, New York, 1978.
- [OCP*06] E. G. Omodeo, D. Cantone, A. Policriti, and J. T. Schwartz. A Computerized Referee. In M. Schaerf and O. Stock, editors, *Reasoning, Action and Interaction in AI Theories and Systems - Essays dedicated to Luigia Carlucci Aiello*, volume 4155 of *Lecture Notes in Artificial Intelligence*, pages 117–139, Springer Berlin/Heidelberg, 2006.
- [OS02] E. G. Omodeo and J. T. Schwartz. A ‘Theory’ mechanism for a proof-verifier based on first-order set theory. In A. Kakas and F. Sadri, editors, *Computational Logic: Logic Programming and Beyond – Essays in honour of Bob Kowalski, Part II*, volume 2408 of *Lecture Notes in Artificial Intelligence*. Springer-Verlag, Berlin, 2002.
- [Urs05] P. Ursino. A generalized small model property for languages which force the infinity. *Matematiche (Catania)*, **LX**(I):93–119, 2005.